

THE DENTAL PRACTITIONER

AND DENTAL RECORD

Including the official reports of the British Society of Periodontology, the British Society for the Study of Orthodontics, the European Orthodontic Society, the Glasgow Odontological Society, the Liverpool and District Odontological Society, the North Staffordshire Society of Dental Surgeons, the Odonto-chirurgical Society of Scotland, and the Dental and Medical Society for the Study of Hypnosis

Editor:

* N. LIVINGSTONE WARD, L.D.S., D.D.S.

Editorial Committee:

- | | |
|---|--|
| * R. V. P. CAMPBELL, F.D.S., H.D.D. R.C.S.
(Edin.), D.D.S. (Amer.), (London) | B. C. LEIGHTON, L.D.S. R.C.S., H.D.D. R.F.P.S.
(King's College Hospital) |
| B. E. D. COOKE, F.D.S. R.C.S., M.R.C.S.,
L.R.C.P. (Guy's Hospital) | Prof. R. W. LOVEL, F.D.S. R.C.S., H.D.D.,
D.D.S. (Newcastle) |
| Prof. A. I. DARLING, M.D.S., F.D.S. R.C.S.,
L.R.C.P., M.R.C.S. (Bristol) | * H. MANDIWALL, M.B., B.S., L.D.S. (Royal
Dental Hospital) |
| Prof. A. D. HITCHIN, M.D.S., F.D.S. R.C.S.
(St. Andrews) | Prof. T. TALMAGE READ, M.Ch.D., F.R.F.P.S.,
F.D.S. R.C.S., H.D.D., L.R.C.P. (Leeds) |
| J. K. HOLT, L.D.S., M.Sc., D.D.S., F.D.S. R.C.S.
(Manchester) | Prof. G. L. ROBERTS, M.B., Ch.B., B.D.S.,
F.D.S. R.C.S. (Sheffield) |
| A. M. HORSNELL, F.D.S. R.C.S., M.R.C.S.,
L.R.C.P. (London Hospital) | * J. E. SEEAR, L.D.S. R.C.S. (London) |
| Prof. A. C. W. HUTCHINSON, D.D.S., M.D.S.,
F.R.S.E. (Edinburgh) | ALAN SHEFFORD, O.B.E., F.D.S. R.C.S.
(University College Hospital) |
| J. IRELAND, L.R.C.P., L.D.S. (Glasgow) | D. S. SHOVELTON, B.Sc., B.D.S., F.D.S. R.C.S.
(Birmingham) |
| I. R. H. KRAMER, M.D.S., L.D.S. R.C.S.
(Eastman Dental Hospital) | * D. F. SOUL, F.D.S. R.C.S. (London) |
| Prof. F. E. LAWTON, B.D.S., D.D.S., F.D.S.
R.C.S. (Liverpool) | Prof. P. J. STOY, B.D.S., F.D.S. R.C.S.
(Belfast) |

* Constitute the Executive Committee

VOL. VIII, No. 12



August, 1958

Monthly 3s. 6d. **JOHN WRIGHT & SONS LTD.** Annually £2 2s.
post 5d. BATH ROAD, BRISTOL 4 post free

SOLE AGENTS: *Australia*: Robertson & Mullens Ltd., Melbourne; *Canada*: The Macmillan Co. of Canada Ltd., Toronto; *New Zealand*: N. M. Peryer Ltd., Christchurch; *Denmark*: Einar Munksgaard, Copenhagen; *Norway*: Olaf Norli, Oslo; *Sweden*: Gumperts Aktiebolag, Göteborg; *South Africa*: P. B. Mayer, Cape Town.



the right setting

The statue of Nelson dominates Trafalgar Square, set off by the fountains, the gardens, and the elegant pillars of the National Gallery.


Human skill dominates the art of dentistry, but to use your skill to its best advantage it must be supplemented by instruments and materials of the finest quality.

Ash products are of the highest possible standard



Head Office & London Showrooms:

AMALCO HOUSE · 26-40 BROADWICK STREET · LONDON · W.1



DENTURAL
DENTURE CLEANSING SOLUTION

SOAKING IS NOT OLD FASHIONED

It still remains the best way to remove tartar, and tea and tobacco stains. It is true that for superficial cleansing there are quicker methods, but some of them are potentially harmful to dentures, and none of them equal soaking in Dentural for the following reasons.

- 1 **Dentural Dissolves Mucin**, the binding agent of tartar. The acid type of cleanser merely dissolves the mineral component, leaving the tough mucin untouched.
- 2 **Dentural Chemical Action Penetrates** between the teeth where the 'rubbing over' process cannot reach.
- 3 **Dentural is Safe** regardless of the strength at which it is used.
- 4 **Dentural will not Initiate Corrosion**. Where metal pins are used for anchoring the teeth the acid type of cleanser can, over a period of time, corrode the metal.

SAVORY & MOORE LIMITED
60-61 WELBECK STREET,
LONDON. W.1



By Appointment to
H.M. The Queen, Chemists

THE DENTAL PRACTITIONER AND DENTAL RECORD

Vol. VIII, No. 12

CONTENTS

August, 1958

	PAGE
EDITORIAL: FLUORIDATION AND ANDOVER	367
THE DIMENSIONAL STABILITY OF THREE SILICONE-BASE IMPRESSION MATERIALS <i>J. N. Anderson, M.D.S., L.D.S. (Sheff.)</i>	368
THE INTEGUMENT OF THE ENAMEL SURFACE OF THE HUMAN TOOTH (<i>continued</i>) <i>E. P. Turner, M.Sc., L.D.S. R.C.S. (Edin.)</i>	373
USE OF AN ALUMINIUM CASTING ALLOY FOR CONSTRUCTING DENTURE BASES <i>D. J. Neill, D.F.C., F.D.S.</i>	382
SILICATE CEMENTS: HOW TO SELECT AND USE THEM <i>George C. Paffenbarger, D.D.S., D.Sc.</i>	387
BOOK REVIEWS	390
ABSTRACTS FROM OTHER JOURNALS	386, 391

ADVERTISEMENTS

Rates: Full page—£15 15s.; Half page—£8 8s.; Quarter page—£4 10s.

Special Positions and Covers: Full page—£17 17s.; Half page—£9 9s.;

Facing text (full pages only)—£18 18s.

Colour—£9 per colour additional to page rate.

Series Discount—10% for 12 insertions, 5% for 6 insertions.

Agency and Publisher's Commission: 10%.

Classified Advertisements: See page xvi and inside back cover.

Type Area: Full page— $5\frac{5}{8} \times 7\frac{5}{8}$ in.; Half page— $5\frac{5}{8} \times 3\frac{3}{4}$ in.; Quarter page— $2\frac{3}{4} \times 3\frac{3}{4}$ in.

Screen for Blocks: Text pages and cover 150.

Copy: First week of preceding month.

For further particulars write to:

Advertisement Manager, John Wright & Sons Ltd., Bath Rd., Bristol, 4 (BRISTOL 75375)

or, for London Area to:

Cecil M. Kaufman, 49, Upper Park Rd., Hampstead, London, N.W.3 (PRIMROSE 0552)

EDITORIAL NOTICES

CONTRIBUTIONS should be sent to the Editor, *The Dental Practitioner and Dental Record*, The Stonebridge Press, Bath Road, Bristol 4, or, if preferred, to any member of the Editorial Committee. Original articles are accepted on the understanding that they are contributed solely to this Journal.

Manuscript should preferably be typewritten with double spacing and wide margins, and the author should keep a copy. Articles and their illustrations become the property of *The Dental Practitioner and Dental Record*, unless authors reserve the right before publication.

Illustrations should be clearly numbered and legends should be written on a separate sheet of paper and not put on the backs of the originals. Each figure should be referred to in the text. Prints are preferred to X-ray negatives and should be on glossy paper. Lettering which is to appear on illustrations is best shown on an overlay or rough sketch. It should not be put on the original.

Tables should be typed on separate pages and each should have a caption which will explain the data without reference to the text.

References to dental literature should be recorded in the text, with the name of the author and the year of publication in parentheses. In the bibliography they should be arranged in alphabetical order in the following form, the abbreviations of periodicals being those adopted in the *World List of Scientific Periodicals* (1952), e.g.:-

SMITH, J. A. K. (1949), *Brit. dent. J.*, **86**, 271.

LEWIS, R. W. B. (1947), *The Jaws and Teeth*, 2nd ed., 471. London: Science Publishing Co.

Authors may obtain reprints of their articles if the Publishers are informed when the proofs are returned. They will be charged at the cost of production and 50 copies are suggested as a minimum.

BUSINESS COMMUNICATIONS should be sent to the Publishers, The Stonebridge Press, Bath Road, Bristol, 4.

Back Numbers.—The Publishers would be glad to purchase copies in good condition of Nos. 1 and 5, Vol. I, and No. 3, Vol. V, of *The Dental Practitioner*.

Binding Cases.—Binding cases for Volumes I to VII may be obtained from the Publishers at 5s., post free. They are supplied attractively lettered in imitation gold on a dark red ground. If desired, the Publishers will undertake the binding at an inclusive charge of 16s. 6d.

REPRINTS

Reprints can be supplied at a reduced price if authors will indicate their requirements at the time that proofs are returned. The following scale will be helpful as a guide to the cost; 50 copies are suggested as the minimum number.

PAGES	NUMBER REQUIRED							
	50	100	150	200	250	300	400	500
2	£ 1 0 0	£ 1 5 0	£ 1 10 0	£ 1 14 0	£ 1 17 0	£ 2 0 0	£ 2 5 0	£ 2 10 0
4	1 15 0	2 4 0	2 13 0	3 0 0	3 5 0	3 10 0	3 19 0	4 8 0
6	2 10 0	3 2 6	3 15 0	4 5 0	4 12 6	5 0 0	5 12 6	6 5 0
8	3 0 0	3 15 0	4 10 0	5 2 0	5 11 0	6 0 0	6 15 0	7 10 0
10	3 10 0	4 7 6	5 5 0	5 19 0	6 9 6	7 0 0	7 17 0	8 15 0
12	4 0 0	5 0 0	6 0 0	6 16 0	7 8 0	8 0 0	9 0 0	10 0 0
14	4 10 0	5 12 6	6 15 0	7 13 0	8 6 6	9 0 0	10 2 6	11 5 0
16	5 0 0	6 5 0	7 10 0	8 10 0	9 5 0	10 0 0	11 5 0	12 10 0
Printed Covers	£ 1 12 6	£ 1 17 6	£ 2 2 6	£ 2 7 6	£ 2 12 6	£ 2 17 6	£ 3 7 6	£ 3 17 6

THE DENTAL PRACTITIONER AND DENTAL RECORD

Vol. VIII, No. 12

August, 1958



EDITORIAL

FLUORIDATION AND ANDOVER

THE announcement of a settlement in the action brought by the Andover Anti-fluoridation Society, who sought a declaration that the practice of putting fluorides in the water-supply was unlawful, is an unfortunate setback for preventive dentistry and preventive medicine as a whole. It must, however, be realized that this was a settlement between the two parties and that this outcome was not decided by a Court of Law. The settlement was achieved by the announcement that the Andover Corporation had withdrawn from the Fluoridation Scheme supported by the Ministry of Health. It seems a tragedy that an important health measure such as this should become a political football on the Local Government stage. Surely such measures should be decided, in the same way as they are controlled, by the Professional Health Authorities or by the Courts of Law. Despite clarification of all doubtful points by a panel of experts at a public meeting in Andover, the anti's have forced the issue, and the newly-elected Corporation has rescinded the decision of the previous Corporation who had made an heroic stand. We sincerely believe that in the long run the people of Andover will regret this action. Superficially it would appear that this Anti-fluoridation Society had a grip on Andover before dental publicity was instigated. Was this a case of poor public relations and of publicity being too little and too late? It would also be interesting to know

what stand was taken by the local dental and medical practitioners on this matter. However, it is all too obvious that in future an intensive propaganda programme must be initiated before such measures are put into effect, and the public must be fully informed of the health reasons beforehand. The Health Authorities of this country do not enter into such measures lightly or without expert advice. It is sincerely to be hoped that other authorities, finding themselves in a similar position, will not consider any such negative action without serious and thorough contemplation of the reasons why fluoridation is advocated. It is only right in a democracy that the local people should be given an opportunity to air their views, but it is doubtful whether they have the right to throw out a measure which has already been implemented without further concrete evidence of any deleterious effects. But for the measures recommended by the experts in preventive medicine and adopted by local authorities, this country would still be riddled with the plagues of the Middle Ages and we would still be suffering from typhoid, small-pox, and cholera. It must always be remembered that in the case of dental caries we are dealing with the most prevalent disease in the world and any measure that can reduce it without disturbing the normal metabolism of the body has a right to be implemented for the good of the community.

THE DIMENSIONAL STABILITY OF THREE SILICONE-BASE IMPRESSION MATERIALS

By J. N. ANDERSON, M.D.S., L.D.S. (Sheff.)

Senior Lecturer in Dental Prosthetics, University of St. Andrews

HYDROCOLLOID impression materials, of both the reversible and irreversible types, show poor dimensional stability on storage due to their high water content. Both types of material lose water to a dry atmosphere and shrink, and also imbibe water and expand if they are placed in a water-bath. Several reports (Hampson, 1955; Skinner, Cooper, and Beck, 1950; Skinner and Pomés, 1946) have appeared dealing with the dimensional changes on storing hydrocolloid impressions, and all draw the conclusion that impressions in these materials should be cast within a short period of time, usually within ten minutes.

Reports on the dimensional stability of the thiokol materials fall into two categories. In clinical tests (Rosensteil, 1955; Pearson, 1955) the change in dimension of an impression during 24 hours does not appear to be of practical significance, though after storage for one week errors of dimension are noticeable. Laboratory testing (Jørgensen, 1957) reveals a contraction both during and after polymerization.

In certain circumstances it may be necessary to store an impression for a period of time; for example, where laboratory facilities are not available near the dentist's surgery. In addition, there are occasions when duplicate models are required. An impression in a material of suitable elasticity and dimensional stability could be cast several times and thus eliminate the necessity for recording a further impression, or duplication of the master model. In addition, it should be noted that the manufacturers of several silicone-base materials recommend that impressions be kept for 6-24 hours before casting them, in order to produce a satisfactory model surface.

MEASUREMENT OF DIMENSIONAL CHANGES

Dimensional changes in elastic impression materials during a period of storage may be

determined in several ways. They may be measured directly by noting the change in the length of a specimen moving freely either in a waxed trough, or, preferably, floating on a mercury bath. The use of a waxed trough to contain the specimen has been applied for many years to determine the setting expansion of dental plasters. Skinner and his associates (Skinner and Kern, 1938; Skinner and Pomés, 1946; Skinner, Cooper, and Beck, 1950) applied this method to the reversible and irreversible hydrocolloids. The mercury bath technique was originally suggested by Docking and Donnison (1948) to measure dimensional changes in investment materials, and later was used by Hampson (1955) and then by Jørgensen (1957) in work on hydrocolloid and thiokol materials respectively.

Criticism may be levelled at these methods that they reveal only the dimensional changes of a freely moving mass of material, and that the results cannot be applied in clinical practice, where the movement of an impression is restricted by the tray.

Other techniques have therefore been used which simulate practical use of the materials. Phillips and Ito (1951), Phillips, Price, and Reinking (1953), and James (1949) recorded impressions of an inlay cavity model, to which a well-fitting inlay had previously been made. The accuracy of fit of this inlay on models made from stored impressions could then be gauged. Phillips and Ito (1951) also noted changes in the relation of markers inserted into the surface of an impression. Skinner, Cooper, and Beck (1950) took impressions of a metal model bearing undercuts, and then made measurements on the resulting stone model after casting the impression. In a later study (Skinner and Hoblit, 1956), the same investigator noted the dimensional changes in impressions taken of an oversize bridge model. The model was one with porcelain teeth in which platinum markers were embedded for reference

points. A round "Invar" peg was used by Thompson (1953) and the resultant plaster model measured by micrometer.

On reviewing the literature, it was felt that one should be able to deduce the direction of

the impression material contracts, the impression cavity should become smaller.

To confirm this deduction for the silicone-base materials, their free dimensional changes were measured, and then the effect of these




Fig. 1.—Dimensional change of three silicone impression materials in 1-7 days.




Fig. 2.—Thermal expansion of three silicone impression materials.

any change in the dimensions of an impression if one knew the changes shown in a freely moving specimen. The direction in which shrinkage or expansion takes place is presumably controlled by the support which the impression material enjoys from the tray. Take the example of an impression within a ring or tube of metal. If the impression material contracts, the impression cavity should increase in size if the material adheres to, or is retained mechanically by, the metal ring. If the impression material expands, the cavity should become smaller. James (1949), Phillips and Ito (1951), and Skinner and Hoblit (1956) commented upon this directional shrinkage for hydrocolloid materials. Conversely, in a completely unsupported impression, if

changes upon supported and unsupported impressions was recorded.

MEASUREMENT OF FREE DIMENSIONAL CHANGES

The use of a mercury-bath for supporting the material gives accurate evidence of the free dimensional changes. Provided that the specific gravity of the material being tested is low, there is little compression or restriction of its movement from the mercury-bath. The apparatus consists essentially of a perspex bath containing mercury. The specimen is attached to one end of the bath, the other end floating freely. Markers are inserted into the specimen and changes in dimension recorded with a comparator microscope.

The materials tested were:—

Lastic 55,
Scania Silicone-base,
Verone.

A weighed mix of the material to be tested was manipulated into a "sausage" shape and placed on the mercury-bath. One end was attached to the end of the bath, and two markers inserted in the floating portion of the specimen. This activity was completed within three minutes of starting to mix. No measurements were taken until the material had set.

Recordings of length were taken at frequent intervals during the first twelve hours, then at one day, and thereafter at daily intervals, until one week had elapsed.

Unfortunately, it was not possible to maintain at 18° C. the temperature of the room in which this experiment was carried out. Over the weekends, when electricity, gas and central heating were cut off, the temperature fell, often to 12° C. Variations in the readings showed a relation to temperature changes, and therefore the mercury-bath temperature was taken at the time of each reading of the microscope.

In order to correct the recorded length of the specimen for temperature changes, it was necessary to determine the material's thermal expansion and contraction. After the 7 days had elapsed, hot water was run into the mercury-bath until it reached a steady temperature of 50° C. The change in the length of the specimen was then noted as the temperature of the bath fell to a low room temperature, usually about 15° C.

The figures for dimensional change of the specimen with time were then corrected for temperature according to the coefficient of thermal expansion which was determined. Recordings of dimensional change were repeated three times for each material and an average result obtained (*Fig. 1*).

All three materials contracted markedly during the first 6 hours, and then the rate of shrinkage was reduced. The figures for contraction after 24 hours are slightly greater than those reported for the thiokol materials (Jørgensen, 1957).

Graphs of the thermal expansion of the three materials appear in *Fig. 2*. From these

the average figures for coefficient of linear expansion (15–40° C.) are:—

Lastic 55	0.00018
Scania S.B.	0.00021
Verone	0.00020

These figures for linear thermal expansion are comparatively high, being twice that given for acrylic resins.

An impression material at room temperature, 18–20° C., will warm approximately to mouth temperature during the 3 or 4 minutes' setting period in the mouth. This will take place rapidly where the material is used in thin sections as the silicone elastomers have good thermal conductivity. After removal from the mouth and cooling to room temperature again, a linear contraction of the order of 0.3 per cent can take place. This contraction is in addition to that shown in *Fig. 1*. The total linear contraction of an impression cast after 24 hours in a plaster mixed with cold tap water could therefore be as high as 0.6 per cent. To reduce the dimensional changes, the impression should be cast in a model material at mouth temperature. This also brings about an improvement in the resulting model surface.

DIMENSIONAL CHANGES IN IMPRESSIONS

In order to discover the practical effects of the supporting medium in controlling the direction of contraction, two experiments were carried out. The first was designed to discover the effect on dimensional changes of the adhesion of silicone-base materials to impression tray materials.

Weighed mixes of each material were placed on pieces of shellac tray material, with and without perforations, and on pieces of stainless steel and nickel-silver. The perforated shellac had 2 mm. diameter holes at 1 cm. spacing. Each support measured 8×3 cm. An impression, 2 mm. thick, was then recorded at 18° C., of a metal ruler bearing centimetre markings. A 5 cm. length appeared on the material over each support. This section was measured by travelling microscope, and changes in dimension noted over a period of three days.

No change in linear dimension was recorded during this period, except with the polished stainless steel support. With this support, however, variable results were obtained, and in only two out of the three materials was a small contraction measured.

At the end of this time, the impression material was carefully removed from its support. In the case of the perforated base-plate, that portion of material passing through the perforations was first cut with a sharp razor. The material was then held at one end of the gauge length and was carefully peeled from the support. Any stretching of the material at this time would, therefore, increase the gauge length. The specimens were placed on a flat surface, and re-measured. A shrinkage smaller in dimension than that recorded in the previous experiment but of wider variation was recorded.

In a second series of experiments, the effects upon the dimensions of an impression were noted.

Impressions of a brass undercut die (Fig. 3) were recorded in one material. The impressions were retained in brass rings and were removed with a sharp pull. Each ring was 0.75 in. in diameter and was perforated with holes of the same size, equidistant from each other.

Six impressions were recorded by this method and after 24 hours they were cast in a weighed mix of dental stone at 18° C. The same batch of stone was used throughout the experiments. One hour after casting, the models were measured across the top and base with a vernier microscope. Measurements were taken along two axes at right angles.

A further series of six impressions was recorded under the same conditions, from the same batch of material. These impressions were carefully removed from the ring by cutting away the material in the retention holes. They were left for 24 hours and then cast and measured as before. Results are given in Table I.

It will readily be seen that those impressions remaining within a ring show an increase in the size of the die, and therefore of the hole within the impression. This is due to a contraction of

the material outwards to its adhesion and retention at the ring. The second group of impressions display a smaller sized die, due to a general contraction inwards of the material.

Experiments in unperforated rings were unsuccessful, as the force required to remove





Fig. 3.—Undercut die used for impressions.

the impression caused the material to pull away from the ring.

Similar results were obtained with all three materials tested. It is considered that the method of measuring the models is not sufficiently accurate to allow correlation of these results with those measured on the much longer, freely movable specimen on the mercury bath.

It would appear, therefore, that if a silicone impression enjoys a good and uniform attachment to a ring or tray, the direction of dimensional change of the impression cavity due to a contraction of the impression material can be

deduced from the examination of the shape of the impression. However, where the material is pulled away from the tray on removing the impression from the mouth, varying dimensional errors can be expected.

silicone materials is higher than that of the thiokols.

The effect on changes in dimension of the adhesion of an impression to its tray or support was also noted. Where adhesion between tray

Table I.—DIMENSIONS OF THE IMPRESSION CAVITY IN SUPPORTED AND UNSUPPORTED SPECIMENS

IMPRESSION	TOP DIAMETER cm.		MEAN ERROR	BASE DIAMETER cm.		MEAN ERROR
	0°	90°		0°	90°	
			<i>Specimens left in the ring</i>			
A1	0.954	0.948	+0.004	0.526	0.532	+0.008
A2	0.948	0.949	+0.002	0.524	0.528	+0.005
A3	0.946	0.948	correct	0.526	0.528	+0.006
A4	0.950	0.952	+0.004	0.530	0.534	+0.011
A5	0.950	0.944	correct	0.529	0.528	+0.008
A6	0.947	0.949	+0.001	0.528	0.531	+0.009
			<i>Specimens removed from the ring</i>			
A7	0.940	0.938	-0.008	0.512	0.508	-0.011
A8	0.942	0.942	-0.005	0.518	0.518	-0.003
A9	0.942	0.940	-0.006	0.518	0.520	-0.002
A10	0.940	0.942	-0.006	0.512	0.518	-0.006
A11	0.940	0.938	-0.008	0.518	0.520	-0.002
A12	0.942	0.942	-0.005	0.512	0.518	-0.006

EFFECT OF HUMIDITY

A considerable amount of literature exists dealing with dimensional changes of the hydrocolloids with alterations in humidity. It was felt that it would be interesting to note any change in dimension of the silicone-based materials under such conditions.

At the end of the 7-day test of free dimensional changes, the specimens were covered with water at 18° C. and left for 24 hours. Very small dimensional changes were noted; generally a very slight expansion took place, instead of a continuation of the contraction recorded during the previous 7 days.

SUMMARY

Three silicone-base impression materials were tested for free dimensional change on storage. All showed a shrinkage slightly greater than that noted for the thiokol materials. The thermal expansion of the

and impression was maintained, the material contracted towards the tray. In an impression of a tooth, this would produce an oversize model. When unsupported, the material contracted to a smaller diameter.

REFERENCES

- DOCKING, A. R., CHONG, M. P., and DONNISON, J. A. (1948), *Austr. J. Dent.*, **52**, 160.
 HAMPSON, E. L. (1955), *Brit. dent. J.*, **99**, 371.
 JAMES, A. G. (1949), *J. dent. Res.*, **28**, 108.
 JØRGENSEN, K. D. (1957), *Acta odont. Scand.*, **14**, 313.
 PEARSON, S. L. (1955), *Brit. dent. J.*, **99**, 72.
 PHILLIPS, R. W., and ITO, B. Y. (1951), *J. Amer. dent. Ass.*, **43**, 1.
 — — PRICE, R. R., and REINKING, R. H. (1953), *Ibid.*, **46**, 393.
 ROSENSTEIL, E. (1955), *Brit. dent. J.*, **98**, 392.
 SKINNER, E. W., COOPER, E. N., and BECK, F. E. (1950), *J. Amer. dent. Ass.*, **40**, 196.
 — — and HOBLIT, N. E. (1956), *J. pros. Dent.*, **6**, 80.
 — — and KERN, W. R. (1938), *J. Amer. dent. Ass.*, **25**, 578.
 — — and POMÉS, C. E. (1946), *Ibid.*, **33**, 1253.
 THOMPSON, M. J. (1953), *Ibid.*, **46**, 1.

THE INTEGUMENT OF THE ENAMEL SURFACE OF THE HUMAN TOOTH

II. THE ACQUIRED ENAMEL CUTICLE

By E. P. TURNER, M.Sc., L.D.S. R.C.S. (Edin.)

Turner Dental School, University of Manchester

(Continued from page 348)

IN Part I of this article the developmental integument of the enamel surface was discussed and it was pointed out that there is disagreement regarding the period of time that the inner layer of Nasmyth's membrane remains on the enamel surface. According to Paul (1896), Stoehr (1910), Lewis (1914), Mummery (1919), Chase (1926), and Atkinson (1950) this membrane is lost shortly after eruption. Choquet (quoted by Dobbs, 1932) considered that it is not present on 88 per cent of erupted teeth and Pincus (1937) states that it remains only in the fissures. The observations of Scott, Kaplan, and Wyckoff (1949), Scott and Wyckoff (1949), and Scott (1952) lend support to these views, by their demonstration of the progressive loss of detail on the enamel surface of erupted teeth as age advances. These authors suggest that this loss is due to wear, and show that a high percentage of the characteristic morphology of the enamel surface of an unerupted tooth is not present within 40 years after eruption, approximately 5 per cent of the loss occurring during the first 10 to 15 years. As would be expected, the loss of detail is more marked in areas of the enamel exposed to wear rather than in the interproximal region, although loss of detail does occur in the latter area, but at a slower rate. However, this evidence has to be reconciled with the fact that a cuticle can be obtained from the enamel surface of an erupted tooth at any age, corroborating the opinion of those, including von Ebner (1922), Walkhoff (1924), Kronfeld (1930), Darling (1943), De Fazio (1947), and Hodson (1949), who consider that Nasmyth's membrane or part of it (e.g., the inner structureless layer or "primary cuticle") remains on the enamel surface throughout life. Approaching the

subject from a different aspect, other writers, including Gottlieb (1921 and 1947), Chase (1926), Kronfeld (1930), von Korff (1930), Dobbs (1932), Bibby and Van Huysen (1933), Manly (1943), Vallotton (1945), Atkinson (1950), Erausquin (1951), Turner (1954 a, b), Rushton (1954), and Waerhaug (1956), describe a structure which forms on the enamel surface (and in some instances on the dentine and cementum if these are exposed to the oral cavity) after a tooth erupts. This cuticle has been referred to by different terms and its origin is disputed. For example, Gottlieb described a keratinous "secondary cuticle" which he stated to be a product of the cells of the epithelial attachment and is referred to by other writers as the "attachment cuticle" (Ussing, 1955). Vallotton (1945) referred to a "brown pellicle" and Campaigne and Fosdick (1938) to "mucin plaques", both cuticles being considered to be derived from the saliva. Kirk (1910) held the view that mucin is deposited from the saliva on the tooth surface by lactic acid producing bacteria. Chase (1926) stated that this cuticle formed as a result of the "vital activities of bacteria". Waerhaug (1956) expressed doubt as to the validity of the view that secondary cuticle formation takes place, and states that "until further evidence has been provided to the proof that there really are two cuticles of different consistency and of different origin, it may be wise to use the unspecific term, enamel cuticle".

MATERIALS AND METHODS

After the fixation of extracted human teeth in 10 per cent formal saline or a fixing agent appropriate to a particular histochemical test, extraneous material was removed by scrubbing

the crown surface with a bristle brush using soap or detergent and tap water.

An obvious principal requirement when obtaining a cuticle to investigate its properties is to separate it from the enamel surface with the minimum risk of altering it histologically or chemically, and not to include the "groove protein" of Pincus (1948). "Physical methods", such as scraping the cuticle from the surface (Manly, 1943; Vallotton, 1945; and Gustafson, 1945) or using an air- or water-jet (Pincus, 1937), were found unsatisfactory as only a small amount of damaged material or no material was obtained. After investigation (although the technique is open to criticism) it became evident that a method of obtaining sufficient quantities of apparently undamaged material was that of Pincus (1937) and Atkinson (1950) in which a tooth is immersed in 2 per cent hydrochloric acid and as the superficial enamel is decalcified the cuticle separates from the tooth surface, this process taking approximately 10 minutes. Even after preliminary cleaning some cuticles were found to have stained extraneous material united with them which rendered the cuticle opaque. Such specimens were discarded.

In view of the observations of Scott, Kaplan, and Wyckoff (1949), and Scott and Wyckoff (1949) which show that some of the structural detail of the enamel surface may be lost during the first ten years after the eruption of a tooth, it is reasonable to assume that Nasmyth's membrane may be altered or lost during this period. In order to deduce any difference in properties between the inner layer of Nasmyth's membrane present on the enamel surface of a recently erupted tooth and the cuticle which is similar in appearance and present on teeth later in life, two groups of specimens were collected.

The first group comprised the inner layer of Nasmyth's membrane (primary cuticle) obtained from partly erupted and recently erupted teeth extracted, for example, during the course of orthodontic treatment. (The method whereby this layer was identified is discussed below.) The second group of cuticles was obtained from non-carious "normal" teeth extracted ten years or more

after eruption, from patients covering a wide range of ages (16-70 years). The appearance and properties of cuticles from each group were then investigated and compared.

To investigate the possibility of the formation of a cuticle on the enamel surface after eruption, the natural cuticle was removed from a posterior tooth extracted from an adult. The roots were cut off and the crown attached by means of a post to a partial denture. This denture was worn in the mouth for periods varying from 7 to 63 days. To accommodate the bite one cusp of the crown was ground, which exposed the underlying dentine. This had the advantage of making it possible to observe whether a cuticle will form on exposed dentine, although it is not suggested that the dentine surface in this instance is identical with attrited dentine.

The cuticles which were found to form on the denture crown and the naturally occurring cuticles of erupted teeth were examined histologically and histochemically and preliminary studies were made using paper partition chromatography.

RESULTS

It was found that a comparison of the inner layer of Nasmyth's membrane with the cuticle present on the enamel surface of teeth erupted for ten or more years showed dissimilarities when the two cuticles are suspended in water and viewed under a low-power microscope. Both cuticles appear on cursory examination to be structureless transparent laminae which may be colourless or lightly pigmented. However, it became evident that certain differences exist between them in appearance and properties as follows:—

1. The inner layer of Nasmyth's membrane exhibits interference phenomena, which is the property of a thin lamina to refract light; in contrast the cuticle removed from a tooth erupted several years does not show this property.

2. When unstained spread preparations are viewed under the high-power objective of a microscope, the inner layer of Nasmyth's membrane invariably shows markings which are usually considered to be due to impressions

formed by outcrops of prisms on the enamel surface (the "perikymata", *Fig. 1*), whereas such markings are faint or non-existent on the cuticle of adult teeth. Frequently in older




Fig. 1.—Unstained spread preparation of the inner layer of Nasmyth's membrane, showing the impressions made by the enamel prism-ends. ($\times 60$.)

teeth it has been found that the latter cuticle is traversed by lines (*Fig. 2*), presenting a similar picture to celloidin replicas prepared from such teeth by Scott and others. It is suggested that these lines are minute "scratches" resulting from wear on the enamel surface which become filled by cuticle material, giving the cuticle this appearance.

3. The two cuticles show different solubilities and different reactions in acid (*see p. 376*).

4. A difference between the cuticles can be demonstrated using histochemical methods (*see p. 377*).

From these observations it is evident that either the inner layer of Nasmyth's membrane alters as age advances or alternatively it is lost and a cuticle forms on the enamel surface after eruption. Both these views have been suggested previously by different authors.

In support of the argument that a cuticle is formed after eruption, it was found that in many cases, when a tooth showing attrition is placed in 2 per cent hydrochloric acid, a cuticle separates from the attrited enamel surface which is continuous with the cuticle covering the remaining surface of the crown. By way of a control it is simple to demonstrate that if an area of the enamel surface is abraded

with a sand-paper disk prior to immersion in acid, a cuticle does not separate from the abraded area. This indicates that a cuticle is formed on the attrited enamel surface, rather




Fig. 2.—Spread preparation of the structureless cuticle removed from the enamel surface of a tooth of a person advanced in age. (Unstained.) ($\times 60$.)




Fig. 3.—Unstained spread preparation of the cuticle formed experimentally on a natural tooth crown. The prism outlines are exaggerated due to etching of the enamel by acid. ($\times 180$.)

than a superficial layer of enamel "dissolving off" in the acid. As a result of this observation experiments were carried out using a natural tooth crown attached to a partial denture. The denture was worn in the mouth for a fixed period of time after which the crown was removed, washed in tap water, and placed in a 2 per cent solution of hydrochloric acid. After several minutes a cuticle separated from the enamel surface as occurs under similar circumstances with an extracted tooth; a cuticle did not separate from the dentine surface. By this method it was found possible to form cuticle after cuticle on the same crown,

the number being limited only by the reduction in enamel area as a result of repeated immersion in the acid used to remove each cuticle. Because the enamel is etched by the acid when the previous cuticle is removed, spread preparations of these "experimental" cuticles showed microscopically the typical morphology of an etched enamel surface (Fig. 3). Under a low-power microscope ($\times 12$) these cuticles resemble the cuticle present on the enamel surface of adult erupted teeth. If the period of wear is less than 12 days, a cuticle separates from the surface which appears normal under the low-power microscope, but is unstable in 2 per cent hydrochloric acid. In this connexion it should be mentioned that the naturally occurring cuticle obtained from areas of enamel attrition sometimes showed a similar reaction. A few minutes after such a cuticle was removed it dissolved slowly in acid. This reaction is discussed below in further detail.

An attempt was made to assess the origin and properties of this cuticle which forms after eruption. For convenience it will be termed the "post-eruption cuticle" in this communication.

THE BEHAVIOUR OF THE CUTICLES TOWARDS INORGANIC SOLVENTS

The reactions of the inner layer of Nasmyth's membrane and the post-eruption cuticle were observed and compared when several cuticles from each group were placed in water at different temperatures and in hydrochloric acid and sodium hydroxide solutions for 24 hours at different temperatures and concentrations which varied from N/100 to 10N. The results are shown in Table I, from which it is evident that the inner layer of Nasmyth's membrane shows a variable reaction, whereas the post-eruption cuticle is insoluble at room temperature in all concentrations of acid. At the same time, the latter cuticle becomes a mauve colour after immersion in the acid. This colour change, together with relative insolubility in acid, serves as a further distinguishing property between the post-eruption cuticle and the inner layer of Nasmyth's membrane.

Generally speaking it can be said that the inner layer of Nasmyth's membrane is insoluble in weak acid, dissolves in strong acid, but between these two extremes the reaction is inconsistent. It was stated in the previous

Table I.—THE REACTION OF THE CUTICLES IN INORGANIC SOLVENTS

HCl	INNER LAYER OF NASMYTH'S MEMBRANE		POST-ERUPTION CUTICLE
10 N to 5 N	Room temp.	Dissolved	Insoluble
	100° C.	Dissolved	Dissolved
5 N to 2.5 N	Room temp.	Dissolved or fragile	Insoluble
	100° C.	Dissolved	Fragile
2.5 N to N/5	Room temp.	Inconsistent reaction	Insoluble
	100° C.	Dissolved or fragile	Fragile
N/50 to N/100	Room temp.	Insoluble	Insoluble
	No colour changes observed.		Colour changes Brown—maroon—mauve

article, in agreement with many writers, that this cuticle appears to be a final layer of enamel matrix. In view of the fact that the cuticle may or may not be calcified this constitution may alter from one tooth to another. Where it is uncalcified it will resemble uncalcified enamel matrix and be relatively insoluble, whereas in teeth where it is partly or completely calcified it will be soluble in acid. This accords with observations on the staining properties of this layer which were described previously.

Neither cuticle showed any change in alkaline solutions at room temperature. At boiling point some specimens became fragile but did not go into solution. Similarly the cuticles did not show any change in water, either at room temperature or at 100° C.

CHEMICAL CONSTITUTION

A biochemical study of the enamel cuticles has been carried out previously by Rosse (1921), Stein, Hinck, and Hoskins (1928), Manly (1943), Vallotton (1943, 1945), and Pincus (1948).

When applying histochemical techniques to examine the post-eruption cuticle it is necessary to exercise meticulous care only to use specimens which appear to be entirely free from extraneous material, otherwise the results are misleading and unreliable.

According to French and Edsall (1945) and Hawk, Oser, and Summerson (1947) formaldehyde can react with a number of functional groups present in proteins. On account of this the ninhydrin reaction was applied directly to fresh unfixed specimens, which gave a negative result. As the specimens were small and thin an alternative method was evolved to apply this test. Several cuticles were hydrolysed in concentrated hydrochloric acid for 15-24 hours at 100° C. The hydrolysate was evaporated to dryness under reduced pressure and the acid was replaced by distilled water. A drop of this solution was absorbed on to a piece of filter-paper and allowed to dry. When the areas were sprayed with a 0.1 per cent alcoholic solution of ninhydrin and the paper warmed over a Bunsen burner flame, a violet colour developed, indicating a positive reaction. A positive result was obtained on applying the Millon and xanthoproteic reaction and the Biuret test, giving evidence of the presence of amino-acids and peptide linkages from which it was concluded that protein is present in this cuticle.

A relatively large volume of humin is formed when several cuticles are hydrolysed in strong acid. Humin is a dark brown insoluble precipitate which forms when a combination of carbohydrate and protein is hydrolysed in acid (Gortner and Blisch, 1915; Roxas, 1916; and Lugg, 1938), this reaction being referred to sometimes as the Maillard reaction. As the presence of protein has been demonstrated already, such a reaction indicates that carbohydrate is also present in the cuticle. This was found to be so, a positive result being

obtained with the thymol test, Benedict's reagent, and Feigl's test, using silver nitrate paper. It has been suggested previously that the post-eruption cuticle is derived from salivary mucin, this substance being a carbohydrate-protein complex.

The literature concerning salivary mucin has been critically surveyed by Clement (1951) and the chemistry of mucopolysaccharides is discussed by Stacey (1946). The behaviour and properties of mucosubstances are described by Kent and Waterhouse (1955).

Histochemical tests were then applied to spread preparations of the cuticle to investigate the combination between carbohydrate and protein shown to be present. The cuticle becomes a red colour when it is stained by the PAS method, indicating the presence of a neutral mucopolysaccharide or mucoprotein. The intensity of the colour produced is some indication of the group to which the tissue belongs (Pearse, 1953). The inner layer of Nasmyth's membrane gives a negative result with this reaction.

Specimens of the post-eruption cuticle and the inner layer of Nasmyth's membrane were fixed in a 16 per cent formalin solution containing 8 per cent lead acetate and stained with toluidine blue using Lison's method (1936). Only the post-eruption cuticle stained metachromatically, becoming a purple-violet colour, indicating a neutral mucopolysaccharide or mucoprotein (Pearse, 1953).

A preliminary study employing paper partition chromatography demonstrated the presence of aspartic acid, glutamic acid, serine, glycine, and alanine. As these amino-acids are present in many proteins this information cannot be regarded as significant. It would be expected that tyrosine would be demonstrated in view of the results obtained with the Millon and xanthoproteic tests. However, it may be that the tyrosine was destroyed during the preparation of the hydrolysate.

The histochemical tests, already described, indicate that the post-eruption cuticle is a mucoprotein or mucopolysaccharide. From this it follows that it may be derived from the mucin of saliva, as suggested by Kirk (1910).

Dobbs (1932), Campaigne and Fosdick (1938), Manly (1943), and Vallotton (1945).

Assuming this thesis to be correct, the question arises as to the mechanism whereby the cuticle is formed on the enamel surface. It was found in a further investigation that an apparently similar cuticle can be formed *in vitro* under sterile conditions on the enamel surface of teeth from which the naturally occurring cuticle has been removed, which

weak acid and examining the cuticles under a dissecting microscope 24 hours later.

It can be seen in sections of fissures and tubular hypoplastic pits that the cells of Nasmyth's membrane degenerate. Debris appears to collect in such areas but invariably a cuticle can be identified covering the enamel surface (Fig. 4). It was not possible to isolate specimens of the cuticle alone from fissures without extraneous material being




Fig. 4.—Section through an occlusal fissure showing part of the cuticle which covers the surface (C). P, Plaque material. D, Debris in the fissure. (H. and E.) ($\times 60$.)

indicates that bacteria do not take part in the process. According to von Korff (1930) mucin is adsorbed on to the enamel surface. Little (1954) states that observations using the electron microscope confirm this point of view. Further work on the problem is necessary before a final conclusion can be obtained.

THE RELATIONSHIP OF THE POST-ERUPTION CUTICLE TO THE ENAMEL SURFACE

The most important connexion of the post-eruption cuticle is its continuity with crack lamellæ (Bodecker and others, 1951; Atkinson and Prophet, 1953). This continuity can be demonstrated readily by immersing a tooth in




Fig. 5.—Section through a plaque (P) on the enamel surface showing the plaque material filling a "wedge" depression (W). E, Enamel. (Gram's stain.) ($\times 300$.)

included. Similarly the cells of Nasmyth's membrane disappear from the wedge-shaped depressions on the enamel surface. Again in such situations a cuticle can be identified. As described by Hodson (1953) these wedge-shaped areas appear to become filled with plaque material (Fig. 5), a fact which may be of significance in early enamel caries.

OTHER SECONDARY CUTICLES

A discussion of the enamel integuments is incomplete if no account is given of the "secondary horny cuticle" originally described by Gottlieb (1921). The existence of such a structure is widely accepted, although some authors doubt this (Hodson, 1949; Waerhaug, 1956). First, it is necessary to indicate how the secondary cuticle can be distinguished from the primary cuticle and the post-eruption cuticle. The most striking difference

is its much greater thickness, which is evident on examining sections of the secondary cuticle. Also this cuticle was found in some instances to contain birefringent substances (Fig. 6).




Fig. 6.—Section of Gottlieb's secondary cuticle (C) viewed by polarized light (using a gypsum plate), showing the birefringent contents which give the cuticle a granular appearance. (H. and E.) ($\times 300$.)




Fig. 7.—Section shows Gottlieb's secondary cuticle in the gingival crevice region with normal nucleated epithelial cells lying adjacent to the cuticle. G, Gingival crevice. E, Space occupied by the enamel. (H. and E.) ($\times 180$.)

The birefringence was usually, but not always, found in that part of the cuticle which was nearest to the oral cavity, and may be due to foreign body material which becomes included in it and may be of significance in early

calculus formation. Where the birefringence occurs the cuticle has a "granular" appearance when viewed in transmitted light. The secondary cuticle was found in sections of erupted human teeth only where the adjacent soft tissue was in situ around a tooth, and then not in all such specimens (although it should be pointed out that Gottlieb inferred that this cuticle is not always present). This cuticle was never identified in sections of cuticles removed from the free enamel surface. However, in longitudinal sections of the secondary cuticle in situ with the epithelial attachment it can usually be seen extending to the cervical region of the clinical crown, which suggests that the cuticle remains confined to the gingival sulcus and its immediate vicinity.

Gottlieb considered the secondary cuticle to be composed of keratin elaborated by the cells of the epithelial attachment. It is difficult to understand how this keratin is produced, as the epithelial cells adjacent to the cuticle appear to be unchanged from the




Fig. 8.—Human. Section showing a laminated cuticle (removed from the gingival crevice area), similar to that described by Rushton as occurring in the hamster. Ep, Epithelial cells from the crevicular epithelium. P, Bacterial plaque. C, Layers of cuticle separated by plaque material. E, Space occupied by the enamel. (H. and E.) ($\times 300$.)

deeper cells and there is a sharp junction between the cells and cuticle in contrast with epidermis, for example (Fig. 7). The thickness

appears regular and the structure amorphous, or granular and striated at right angles to its length, unlike keratin layers elsewhere which are often irregular in thickness and somewhat "scaly" in appearance. Furthermore, the cuticle does not readily stain or react similarly to keratin with commonly used stains such as haematoxylin and eosin and Mallory's stain. Quite apart from this evidence it is generally accepted that physical stimulation of an epithelial surface is an important factor in keratin formation, and it is doubtful if such stimulation exists in the situation where the secondary cuticle is found. Trott (1957) did not find evidence of keratinization of this crevicular epithelium in his study of the gingival epithelium.

The acquired enamel cuticle described by Rushton (1954) as occurring in hamsters must also be mentioned. Such a thick, laminated, yellowish, non-birefringent cuticle can be seen associated with human teeth (*Fig. 8*), but its presence appears to be confined to stagnation areas where there is plaque material and to carious teeth. As this type of cuticle has been found to occur only in the pre-carious or carious state it was considered that it could be included better in a study of the significance of the cuticles in relation to caries, and therefore it was not included to any extent in the present study. The areas covered by this cuticle are relatively small and it has a segmented appearance in cross-section.

Both Gottlieb's secondary cuticle and that described by Rushton give a positive result with a modified Gomori's aldehyde fuchsin method. As a control, sections of mucous gland and keratinized oral mucous membrane were stained under identical conditions using the same batch of solutions; the former gave a positive result and the latter was negative, which demonstrates that the cuticles under discussion are not composed of keratin and may well be mucinous in origin. However, it was found that they do not show the reaction given by mucinous material when stained by Schiff's method, or with Toluidine blue, Celestin blue, Alcian blue, and Mucicarmine (sections of gastric mucous tissue being used as a control in each case). It is hoped to

explain this contradictory behaviour after investigation of age changes in the cuticles.

In human teeth both cuticles are present, mainly in sheltered areas rather than the free enamel surface and in intimate association with other structures (epithelium or plaque), which adds difficulty to their investigation as they cannot be isolated for analysis in the way that was possible with the primary cuticle and post-eruption cuticle. Further work on these cuticles is in progress.

DISCUSSION

Although the work described is concerned with the integument of the "free" enamel surface rather than the fissures, it is appreciated that the latter are of first importance when considering dental caries. However, it was felt that an attempt should be made to clarify in general terms the conflict of opinion regarding the formation and nature of the enamel cuticles. Authors differ in their views of the significance of the cuticles when considering dental caries or the solubility of surface enamel in acid; some even ignore them.

The mechanism whereby the post-eruption cuticle forms is not understood. It is not apparently the result of a simple deposition of mucin because the cuticle is often found to be relatively soluble in acid where it is removed from areas of constant wear, but very resistant when obtained from the "free" enamel surface. This resistance cannot be explained by suggesting that on the "free" surface the cuticle becomes greatly increased in thickness because this does not appear to be so and there is a limit to the thickness attained. The change appears to be in the composition of the cuticle itself.

In spite of the conflicting results obtained in a preliminary examination of Gottlieb's secondary cuticle and the laminated cuticle similar to that described by Rushton, the writer is inclined to the view that the secondary cuticle and the laminated cuticle originate from the same source as the post-eruption cuticle and that all three cuticles are a modification of the same structure. In this connexion it is interesting to note that following the mechanical removal of the

secondary cuticle from the gingival pockets of experimental animals, Waerhaug (1956) found that on subsequent investigation the cuticle re-formed after a period of time. From this Waerhaug concluded that the cells of the epithelial cuff re-formed the cuticle. This observation could also be interpreted to suggest that the cuticle was formed in the same way as that on a denture crown experimentally, as described above. That this cuticle and the laminated cuticle are thicker could be explained by the fact that both cuticles form in sheltered areas, which accords with the observation already discussed that the thick secondary cuticle is not found on the free enamel surface, except in the gingival region.

A study of the literature describing the protection and resistance of the enamel surface to caries or acid solution gives an indication of the possible significance of the post-eruption cuticle. Of particular interest in this respect is the work of Darling (1943), who found experimentally that the enamel of erupted teeth possesses a cortical protection against artificial caries and dilute acid except where there is natural or artificial abrasion or attrition; this observation accords with the writer's finding described above that the post-eruption cuticle recovered from such areas is relatively soluble in acid. Walsh and Green (1950), who demonstrated that the enamel surface can be protected by an adsorbed film of a long chain amine, observed that prior to their experiments the application of sticky wax to the enamel surface and its removal with carbon tetrachloride in some way alters the solubility of the surface enamel in acid attack. It is tempting to suggest that the post-eruption cuticle was removed in the process of applying and dissolving off the sticky wax, thus allowing the enamel to be more readily dissolved by the acid.

It has been demonstrated by Manley and Hardwick (1952) that in enamel caries the affected enamel matrix is rendered resistant to acid, and these authors state that "mainly muco-protein" from the saliva is "absorbed onto" the porous enamel. It follows that such substances may be similar to those composing

the "post-eruption cuticle" and may also occupy faults in the enamel to form "crack" lamellæ (Atkinson and Prophet, 1953).

SUMMARY

Evidence is presented in this and the previous communication indicating that Nasmyth's membrane is composed of an outer cellular layer which is removed by the trauma of mastication shortly after eruption and an inner structureless layer which remains for several years after eruption. The presence is demonstrated of a cuticle which appears to be derived from saliva, and termed the post-eruption cuticle. A study of the composition and properties of these cuticles is described.

Acknowledgements.—The writer wishes to thank Professor R. Dennell, Professor of Experimental Zoology, University of Manchester, for his invaluable advice and assistance and for providing the facilities of his department where much of the work was carried out. Also Professor H. G. Radden for his interest in the work and the histology laboratory technicians of the Turner Dental School for their assistance.

REFERENCES

- ATKINSON, H. F. (1950), D.D.S. Thesis, University of Manchester.
 — — and PROPHET, A. S. (1953), *Brit. dent. J.*, **95**, 60.
 BIBBY, B. G., and VAN HUYSEN, G. (1933), *J. Amer. dent. Ass.*, **20**, 828.
 BODECKER, C. F., GOTTLIEB, B., ORBAN, B., ROBINSON, H. B. G., SCHONS, I., and SOGNAES, R. F. (1951), *Oral Surg.*, **4**, 787.
 CAMPAIGNE, E. E., and FOSDICK, L. S. (1938), *Northw. Univ. Bull. dent. Res.*, **38**, 14.
 CHASE, S. W. (1926), *Anat. Rec.*, **5**, 357.
 CLEMENT, A. J. (1951), *J. dent. Ass. S. Afr.*, **6**, 277.
 DARLING, A. I. (1943), *Proc. R. Soc. Med.*, **36**, 499.
 DOBBS, E. C. (1932), *J. dent. Res.*, **12**, 581 and 853.
 VON EBNER, V. (1922), *Handbuch der Zahnheilkunde*, **1**, 325.
 ERAUSQUIN, J. (1951), *Int. dent. J., Lond.*, **1**, 10.
 DE FAZIO, E. (1947), *La Stomatologia*, **2**, 245.
 FRENCH, D., and EDSALL, J. T. (1945), *Advances in Protein Chemistry*. New York: Interscience Publications.
 GORTNER, L., and BLISCH, J. (1915), *J. Amer. chem. Soc.*, **37**, 1630.
 GOTTLIEB, B. (1921), *Dtsch. Mschr. Zahnheilk.*, **39**, 142.
 — — (1947), *Dental Caries*. London: Kimpton.
 GUSTAFSON, G. (1945), *Odont. Tidskr.*, **53**, supplement.
 HAWK, P. B., OSER, B. L., and SUMMERSON, W. H. (1947), *Practical Physiological Chemistry*, 12th ed. London: Churchill.

- HODSON, J. J. (1949), *Brit. dent. J.*, **87**, 167.
 — (1953), *Ibid.*, **94**, 141.
 KENT, P. W., and WATERHOUSE, M. W. (1955), *Biochemistry of the Aminosugars*. London: Butterworth.
 KIRK, E. C. (1910), *Dent. Cosmos*, **52**, 729.
 VON KORFF, K. (1930), *Dtsch. Mschr. Zahnheilk.*, **11**, 817.
 KRONFELD, R. (1930), *J. Amer. dent. Ass.*, **17**, 1889.
 LEWIS, F. T. (1914), *Textbook of Histology*. Philadelphia: Blakiston.
 LISON, L. (1936), *Histochemie Animale*. Paris: Villars.
 LITTLE, K. (1954), Personal communication.
 LUGG, J. W. H. (1938), *Biochem. J.*, **32**, 775.
 MANLEY, E. B., and HARDWICK, J. L. (1952), *Brit. dent. J.*, **92**, 225.
 MANLY, R. S. (1943), *J. dent. Res.*, **22**, 479.
 MUMMERY, J. H. (1919), *Phil. Trans.*, **209**, 304.
 PAUL, F. T. (1896), *Brit. dent. J.*, **37**, 1021.
 PEARSE, EVERSON (1953), *Histochemistry. Theoretical and Applied*. London: Churchill.
 PINCUS, P. (1937), *Brit. dent. J.*, **63**, 139.
 — (1948), *Biochem. J.*, **42**, 219.
 ROSSE, W. (1921), *Z. Stomat.*, **19**, 348.
 ROXAS, L. (1916), *J. biol. Chem.*, **27**, 71.
 RUSHTON, M. A. (1954), *Brit. dent. J.*, **97**, 199.
 SCOTT, D. B. (1952), *Oral Surg.*, **5**, 638.
 — KAPLAN, H., and WYCKOFF, R. W. G. (1949), *J. dent. Res.*, **28**, 31.
 — and WYCKOFF, R. W. G. (1949), *J. Amer. dent. Ass.*, **39**, 275.
 STACEY, M. (1956), *Advanc. Carbohydr. Chem.*, **2**, 161.
 New York: Acad. Press Inc.
 STEIN, J. B., HINCK, C. F., and HOSKINS, M. M. (1928), *Dent. Cosmos*, **70**, 592.
 STOEHR, P. (1910), *Histology*. Philadelphia: Blakiston.
 TROTT, J. R. (1957), *Brit. dent. J.*, **103**, 421.
 TURNER, E. P. (1954 a), *J. dent. Res.*, **33**, 732.
 — (1954 b), *Nature, Lond.*, **173**, 596.
 USSING, M. J. (1955), *Acta odont. Scand.*, **13**, 123.
 VALLOTTON, C. F. (1943), *J. dent. Res.*, **22**, 204.
 — (1945), *Ibid.*, **24**, 161, 171, and 183.
 WAERHAUG, J. (1956), *Ibid.*, **35**, 313.
 WALKHOFF, O. (1924), *Die Normale Histologie Menschlicher Zähne 2te*. Leipzig: Felix.
 WALSH, J. P., and GREEN, J. W. (1950), *J. dent. Res.*, **28**, 270.

USE OF AN ALUMINIUM CASTING ALLOY FOR CONSTRUCTING DENTURE BASES

By D. J. NEILL, D.F.C., F.D.S.

Department of Dental Prosthetics, Guy's Hospital Dental School

ALUMINIUM is obtained by electrolytic reduction from its oxides and silicates and the resultant metal is a commercial grade of aluminium containing up to 1 per cent of impurities. By a further electrolytic refining process the metal may be obtained with only 0.01 per cent of impurities and in this form is known as "super purity aluminium". This is very resistant to corrosion and is capable of producing a highly reflective surface but it is soft and lacking in strength. These properties may to some extent be modified by alloying with other elements.

The use of "super purity aluminium" for construction of bases for full upper dentures has been advocated on several occasions. Campbell (1935) described a technique in which the material was first cast and subsequently swaged and the following year described the use of an alloy of super purity aluminium with 3 per cent magnesium for construction of bases by casting alone. Sizeland-Coe (1951) advocated the use of a similar alloy in the wrought form by swaging or pressing and suggested that this material was suitable for

limited use as a partial denture base material. Tregarthen (1944, 1949) who used both the casting and swaging methods did not state the type of alloy he used but he described in detail the method of anodizing the prepared base.

Whilst the most enthusiastic advocate of the use of aluminium alloys could not claim that they are ideal as materials for the construction of denture bases, there are occasions when they may be employed for this purpose with advantage.

Since many practitioners may be unfamiliar with the use of these alloys it is proposed to give details of a suitable casting alloy and after comparing its properties with those of other denture-base materials to discuss indications for its use. This will be followed by a consideration of the particular problems associated with the casting of this type of alloy and details of the method employed in processing and finishing denture bases in this material.

The metal which has been used successfully in this department over the past two years is an alloy, not of the super purity type, but of

the commercial grade of aluminium containing 5 per cent magnesium and conforming to the specification LM 5. Its cost is approximately 1½d. oz. which compensates to some degree for the expense involved in fabricating a metal base.

strong to permit their use in a skeleton construction they are strong enough to be used in the form of plates.

There are, however, two outstanding objections to the use of these materials in the partially edentulous case.

Table I

	ALUMINIUM ALLOY	ACRYLIC RESIN	CHROME-COBALT ALLOY	GOLD ALLOY
Density	2.66	1.18	8.2-8.6	15
Hardness (Brinell)	60-68	23-29	280	softened 138 hardened 210
Ultimate tensile strength (tons per sq. in.)	9.6	3	49	softened 26 hardened 49
Melting range (°C.)	580-640	—	1270-1305	870-985
Percentage elongation	1076 40 F 1184	—	5	softened 4-25 hardened 1-6

Table I shows the approximate physical properties of this alloy together with those of other denture base materials for comparison.

Whilst it will be seen that the aluminium alloy is inferior in strength to the other metallic bases it is three times stronger than acrylic resin and if used only for plate types of construction gives excellent results.

In hardness it is again inferior to the other metals used in denture construction but it is more than twice as hard as acrylic in the cast state, takes a higher polish, and irregularities can be removed from the fit surface without destroying its close adaptation to the tissues. Both strength and hardness are increased by the anodic film which may be imparted to this material.

With regard to weight the aluminium alloy is by far the lightest of the metallic bases. Allowing for the thinner section of the palate a denture cast in this material approximates closely in weight to one constructed entirely of acrylic resin.

In addition this alloy has a high thermal conductivity (0.4 c.g.s.) and is resistant to corrosion in the anodized state (see p. 384).

INDICATIONS FOR THE USE OF THE CAST-ALUMINIUM ALLOY

This material is capable of fabrication by casting into partial dentures of limited design. Although these alloys are not sufficiently

1. The possibility of amalgamation between the aluminium of the base and the mercury from new dental restorations. This affinity of



Fig. 1.—A defective casting caused through failure to use a satisfactory flux.

aluminium for mercury results in a peculiar filamentous growth of aluminium oxide on the metal.

2. Electro-chemical corrosion of the base material in the presence of gold and amalgam fillings. Whilst anodizing affords a high degree of protection against both chemical and electro-chemical dissolution the smallest defect in this oxide layer results in quite a considerable electrical potential being developed

between the exposed metal and any gold or amalgam restoration with which it comes into contact.

This can be readily demonstrated by placing a specimen of wrought gold wire and a strip of anodized aluminium in contact with each other in a solution of artificial saliva and connecting them together through a high-impedance volt meter. If the aluminium

where a metal base is indicated but retentive factors are unfavourable.

PROBLEMS ASSOCIATED WITH CASTING AND DETAILS OF CASTING TECHNIQUE

The casting of aluminium alloys, particularly those containing high proportions of magnesium, calls for special care. Mishandling of the material and faulty technique may result in gross casting defects (*Fig. 1*). In addition to the common oxidation difficulties there is the problem of molten gas inclusion, particularly hydrogen, which may be liberated by the reaction of the magnesium in the alloy with any residual water contained in the refractory crucible. Aluminium alloys are also susceptible to grain growth.

To deal with these problems low-fusing fluxes comprising mainly fluorides and chlorides of magnesium and potassium are used and a grain-refining agent is employed which also cleanses the molten metal of metallic oxides and non-metallic inclusions.

Because of the fluidity of the low-fusing fluxes care must be taken to prevent them being included in the metal during casting. This is accomplished by the use of a "drossing-off" compound—a dry flux which forms a skin on the surface of the molten metal and is applied just before casting.*

Although the co-efficient of expansion of this alloy is high its melting temperature is lower than that of gold and it has been found that accurate castings may be obtained using a proprietary gypsum bound cristobalite refractory.

After the master model has been prepared by the incorporation of a post-dam area and by relieving when and where necessary, a duplicate model is prepared in this refractory and a wax pattern of the base is laid down. The thickness of the wax may be varied according to the strength required in the final casting. For a flat palate one layer each of No. 8 and No. 5 waxes may be used, whilst a V-shaped palate with well-pronounced rugæ




Fig. 2.—Pattern waxed-up on investment model showing the distribution of sprues (a riser is attached at the centre of the palate to improve the flow of the molten metal).

specimen has been well anodized no potential difference will be recorded but if now the surface layer of this specimen is stoned away and the electrodes are brought together again a marked potential difference will be recorded.

Minor deficiencies in the protective anodic film will render the alloy more susceptible to attack by chemical agents but under normal mouth conditions the base material remains unaffected by corrosion even when it has been necessary to destroy part of the oxide layer in easing a denture.

In view of the foregoing it is necessary to restrict the use of aluminium alloys to edentulous mouths, and they are used to best advantage in the construction of full upper dentures

* The fluxes used are supplied by Foundry Services Ltd., and will be referred to by their proprietary names in the description of the casting process.

will probably be strong enough if cast in a single thickness of No. 8 wax. Retention tags and a palatal finishing line are added and the case is then ready for spruing; the sprues are formed from round wax of 2.5 mm. diameter and these are attached as shown in Fig. 2.

Investment is completed in a suitable-sized casting-ring, and, after drying out, the mould is heated to eliminate the wax in the usual manner.

For dental purposes it is necessary to use a centrifugal casting technique as in both air- and steam-pressure methods gas inclusion is likely to occur.

In view of the low specific gravity of the alloy it is necessary to employ a high initial velocity to the centrifugal machine to achieve a satisfactory casting pressure and so, prior to heating the mould, the casting machine should be suitably adjusted.

The heated investment mould should be removed from the furnace and placed into position on the carriage of the casting machine whilst the sprue hole is glowing red. This allows adequate time for the mould to cool to its correct temperature whilst the alloy is being prepared.

A dry crucible is charged with an ingot of the alloy approximately 45 g. in weight and this is heated in an open furnace. Before fusion takes place the surface of the metal is covered with a powdered flux known as Coveral 65. The metal is now gently heated with the reducing flame of a gas-air blow-pipe in order to get the mass fluid. Care should be taken not to disturb the metal surface. (Unlike gold alloys this metal should not be spun.)

Into this molten metal is now plunged 0.5 g. of the grain refining agent (Degaser 190) and this is held in place until all reaction is complete. This is followed by the addition of the drossing-off compound (Coveral 11), which should be sprinkled over the surface of the melt and stirred into the metal. The crucible containing the molten metal should now be carefully transferred to the casting machine and when in place casting should be effected without delay.

After casting, the ring and its contents should be bench-cooled before dis-investing.

MECHANICAL FINISHING AND ANODIZING OF THE CASTING

One of the sprues is left when the plate is separated from the remainder of the casting in order that the base may be totally immersed in the cleansing and sealing solutions and that a




Fig. 3.—Diagrams of the circuit of the anodizing apparatus. A, Using mains current supply; B, Using a 12-volt battery.

satisfactory electrical contact may be made between the metal and the anode clamp outside the electrolyte.

After shot-blasting the metal surface is mechanically finished by means of stones and polished with pumice and whitening and metal polish. The polished base is now rinsed under a running tap and scrubbed with a soft brush and detergent powder to remove all traces of abrasive materials. It is next immersed in carbon tetrachloride for 2 minutes to remove any grease attaching to it. Between this and each of the subsequent chemical immersions the specimen should be rinsed in distilled water.

In order to render the surface chemically clean and thus susceptible to electrolytic activity, the base is immersed in a 5 per cent solution of sodium hydroxide for one minute and any excess of the alkali remaining is neutralized by rinsing in 10 per cent nitric acid for a further minute.

The specimen is now made the anode in an electrolytic bath in which the electrolyte is a 10 per cent solution of sulphuric acid and the cathode is formed by a pair of lead or carbon rods.

A direct current is supplied through a combined transformer-rectifier (or alternatively a 6-volt or 12-volt car battery may be used) and this is adjusted so that the current density is about 50 milliamperes per square inch. At this rate the anodizing process takes 45 minutes.




Fig. 4.—A completed case.

The solution is initially at room temperature and there is a small rise in the temperature of the electrolyte which occurs during anodizing. The apparatus used is essentially the same as that commonly employed for copper-plating inlay impressions, and the circuit diagrams are shown for both mains and battery activated equipment (Fig. 3).

When this process is complete the specimen is removed from the circuit, washed and boiled in distilled water for 10–15 minutes in order to seal the protective layer. At this stage the remaining sprue is removed and the base plate may be handled.

After the teeth have been waxed up on the aluminium base and "tried-in" satisfactorily, the case is flaked and processing of the

acrylic resin gum work is carried out in the usual way.

After the finished case is removed from the plaster, care must be taken to ensure that the anodized surface is not damaged whilst polishing the acrylic surface. (Fig. 4 shows a completed case.)

SUMMARY

The use of a commercial alloy of aluminium containing 5 per cent of magnesium for the construction of denture bases by a centrifugal casting technique is discussed.

The properties of this alloy are compared with those of other denture base materials and indications for its use are given.

The technical problems involved in the handling of this material are considered and details of the method of construction and the finishing of aluminium denture bases are presented.

Acknowledgements.—I wish to thank Professor H. B. Fenn for his encouragement and advice in the preparation of this paper. I am indebted to Mr. J. H. Cragg for the part he has played in developing the laboratory technique and to the British Aluminium Co. Ltd., and Foundry Services Ltd., for their technical assistance.

In conclusion I wish to express my gratitude to Miss B. Whiteley of the Department of Dental Photography and Miss D. Carter of the Department of Medical Illustration for the photographs and diagrams.

REFERENCES

- CAMPBELL, D. D. (1935), *J. Amer. dent. Ass.*, **22**, 2082.
 — (1936), *Ibid.*, **23**, 1264.
 SIZELAND-COE, J. W. (1951), *Brit. dent. J.*, **91**, 267.
 TREGARTHEN, G. G. T. (1944), *Ibid.*, **76**, 300.
 — (1949), *Ibid.*, **86**, 227.

Vinyl Ether

Although vinyl ether is extremely volatile and therefore difficult to give, there is a very high degree of safety as against ether or ethyl chloride administered with nitrous oxide and oxygen.

The disadvantages are that it decomposes rapidly when exposed to air, it is expensive, and it has a garlic-like smell which is very unpleasant. In spite of its disadvantages the safety in administration is very great, and there are no complications.—MASSEY DAWKINS, C. J. (1958), *Brit. med. J.*, **1**, 1116.

SILICATE CEMENTS: HOW TO SELECT AND USE THEM*

By GEORGE C. PAFFENBARGER, D.D.S., D.Sc., Washington, D.C.

DENTISTS know the shortcomings of silicate cement. The high solubility, low strength, extreme brittleness, high shrinkage, and acidic and irritating character of the cement are tolerated simply because no other material has a better combination of desirable attributes as a direct filling material for the front teeth. Therefore, the method of using silicate cement is all-important. The values for solubility, strength, shrinkage, and the amount of acidic products available for tissue irritation can be varied depending upon the method of using the cement. The demonstration of these effects in the laboratory and the transfer of these laboratory findings to clinical practice are the primary purpose of this report.

Composition and Its Effects

Powder.—The powders are complex aluminosilicates, usually fused with a fluoride containing flux (Matthews, 1935; Paffenbarger, Schoonover, and Souder, 1938). The fluorine content is relatively high—ranging as high as 15 per cent by weight (Paffenbarger, Schoonover, and Souder). This high fluorine content causes a silicate cement to etch glass. It probably accounts for the comparatively small amount of recurrent caries about silicate cement restorations (Volker, Bekaris, and Melillo, 1944) as silicate cement produces a considerable reduction in enamel solubility (Phillips and Swartz, 1957).

The glass-like powder is relatively insoluble in distilled water. In fact, it is about one half as soluble as powdered enamel in the same medium (Paffenbarger, Schoonover, and Souder).

Liquid.—The silicate cement liquids are aqueous solutions of phosphoric acid partially neutralized by the addition of aluminium or zinc salts or both (Paffenbarger, Schoonover, and Souder). The free phosphoric acid content

will run from 35 to 47 per cent and have a pH of approximately 2. There is no such thing as a "neutralized" silicate cement liquid. Because of the high phosphoric acid content, the silicate cement liquids gain or lose water depending on the humidity of the air to which they are exposed. This hygroscopic property of phosphoric acid is one of the factors that make the relative humidity of the air in which the cement is mixed of some importance.

Care of Powder and Liquid

The powder is stable in almost any climate. The chief precaution is to keep the bottle of powder tightly stoppered at all times, except when removing the powder for mixing. Slight contamination will make a perceptible colour change.

The liquid being susceptible to loss or gain of water should be protected from the air as much as possible. This means that (1) the stopper or cap should be in the bottle at all times except when actually withdrawing the liquid; (2) the liquid should not be placed on the glass mixing slab until just prior to beginning the mix; (3) the smallest possible area of the slab should be used in making the mix; and (4) ideally, the mixing should be done in a container which would protect the mix from the atmosphere (Grunewald, Dickson, Paffenbarger, and Schoonover, 1953).

Two methods of protecting the cement liquid from gain or loss of water due to atmospheric conditions are shown in Fig. 1. The saturated salt solution in the desiccating jar on the left must have a vapour pressure nearly the same as the silicate cement liquid. This will vary with the compositions of the liquid, so inquiry should be made to the manufacturer of the particular trade brand being used concerning the particular air humidity at which the vapour pressure of the cement liquid is in equilibrium. Different salts must be used depending upon

* A paper delivered to the American Dental Society of Europe, August 28, 1957.

the relative humidity desired. Standard chemical handbooks list the relative humidity over many salt solutions (*Handbook of Chemistry and Physics*). To maintain saturated solutions, there must always be some undissolved salt in the bottom of the desiccating jar.

Another suggestion is shown on the right of Fig. 1. Light liquid petrolatum, U.S.P., is added to the cement liquid. This spreads out




Fig. 1.—Two methods of protecting the cement liquid from loss or gain of water due to atmospheric conditions.

into a thin film as shown in the illustration and effectively prevents changes in the water-content of the liquid.

Microstructure of the Hardened Cement

Silicate cement powder is slowly dissolved in the cement liquid and in any practicable mix there is always powder in excess. Consequently, the hardened silicate cement appears in cross-section as shown in Fig. 2. The outline of the shape of the original powder particles can be seen. The cement liquid only attacked the surfaces of the larger particles. The product of this reaction between the cement liquid and the surface of the particles of powder is called the matrix. It is the substance that binds the undissolved powder particles together. It is the most soluble and weakly acidic portion of the hardened cement. The matrix is the part which shrinks, stains, and loses water in contact with air. Thus, the gel-like colloidal matrix, or the medium that cements the unused powder particles together, is the part

that gives the silicate cement its undesirable properties. It is therefore important that the amount of matrix present in a mix of hardened cement should be kept minimal. This can be accomplished by a proper mixing technique.




Fig. 2.—Cross-section of a hardened silicate cement. (The dark areas are bubbles of air, one defect caused by improper mixing.)

Technique of Mixing

Cold Slab.—The mixing slab should be chilled to just above its dew point. This retards the reaction between the powder and liquid during mixing. Thus, a larger amount of powder can be incorporated in a given quantity of liquid when the mix is made on a chilled rather than on a warm slab. If one makes the mix in a small rubber sack under cool water, then the maximum amount of powder can be incorporated regardless of atmospheric conditions.

Also, the mix should be made as rapidly as possible to minimize the formation of excess matrix. A mix can be made in 30 sec. with just a little practice. Longer mixing times increase solubility. The purpose of the mixing is to wet completely the powder surface with the liquid and to start the reaction between powder and liquid. As small an area of the slab as is practicable should be used to minimize exposing the hygroscopic unset

cement to the atmosphere. The mix should attain a heavy putty-like consistency. In short, one should make a thick mix on a cool slab as fast as possible. This will ensure the minimum amount of the matrix material which contributes to the most undesirable properties.

The dark circular areas in *Fig. 2* are air-bubbles entrapped during the mixing of the cement.

Technique of Insertion

As soon as the mix is completed, it should be transferred to the cavity and the retaining or contouring strip should immediately be placed over it, drawn taut and held unvarying in this position until the cement has hardened. Do not guess at this time of hardening. Make the mix in excess and after insertion of the restoration hold the excess in the palm of one hand. Test its relative hardness by indenting with the finger-nail. When the cement cannot be indented, it has set hard enough to remove the strip. Make every attempt to judge the correct amount of cement to pack into the cavity so that the minimum amount of finishing will be required. After the strip is withdrawn, the filling should be coated with a grease such as petrolatum U.S.P., or preferably a silicone stopcock grease such as is used in technical laboratories. This grease is very tenacious and water repellent. The purpose of the coating is to keep the filling protected from saliva or air. In other words, the coating will preserve the water-content of the filling for a few hours after insertion. Some dentists prefer a low-melting paraffin wax as a coating. Wax is indeed an effective water barrier.

Technique of Polishing

None of the abrasives in the dental office will produce a surface as smooth as that left by the retaining strip. The lingual surface should require little or no finishing if the strip has been properly adjusted.

Polishing should be delayed for at least 24 hrs. as the cement is very weak just after the removal of the retaining strip. The data in *Table I* show how the strength of a silicate cement changes with time. The finest grit of cuttle-fish disks should be used slowly and should be covered with a grease so that the

surface of the filling will be protected from the air. Under no circumstances should the filling be made flush with the enamel margin just after removal of the strip. To do so breaks off

Table I.—CHANGE OF COMPRESSIVE STRENGTH OF A SILICATE CEMENT WITH TIME

Age of Specimens Time	Compressive Strength lb. per sq. in.
15 min.	10,000
1 hr.	12,000
3 hr.	15,000
1 d.	16,000
1 wk.	19,500
6 m.	22,500
14 m.	24,000

the edges of the weak silicate cement and makes a V-shaped ditch around the filling. Much of the damage to silicate cements is done by polishing prematurely.

Care of Silicate Cement Filling

If the patient is a mouth breather, he should be instructed to coat his silicate fillings with petrolatum every night before retiring; otherwise the silicate cement restorations will last only a few months. The dentist should always protect any silicate cement fillings that he may isolate with rubber dam or cotton rolls by coating them with a grease, preferably a silicone stopcock grease.

Demonstration of the Effect of Technique upon Fillings in the Mouth

Laboratory data on silicate cements demonstrate the effect of technique upon the numerical values for pertinent chemical and physical properties such as setting time, solubility, strength, and shrinkage. These effects can likewise be demonstrated clinically.

In *Fig. 3* are two adjacent silicate cement fillings made of the same cement, which show gross differences within even a few weeks of insertion (Paffenbarger, 1951). The good filling on the left was made from a mix spatulated for 30 sec. on a slab at 60° F. Petrolatum was placed on this filling just after the matrix strip was removed. Finishing was deferred for one week. The restoration on the right was made from a mix spatulated for 2 min. on a slab at 98° F. Moisture came in contact with the restoration after the matrix strip was removed.

Immediately thereafter, the restoration was dressed down flush with the enamel margins.




Fig. 3.—Silicate cement fillings made from the same cement but by proper techniques (left filling) and improper techniques (right filling).

The photograph (Fig. 3) was taken a few weeks after the fillings were inserted.

Conclusion

A silicate cement restoration may be either a work of art or a disappointment to the

patient and the dentist. The difference can depend on the technique used in mixing, packing, and finishing the restoration. This can be demonstrated readily on controlled clinical tests constructed on the basis of data derived in the testing laboratory.

REFERENCES

- GRUNEWALD, A. H., DICKSON, GEORGE, PAFFENBARGER, G. C., and SCHOONOVER, I. C. (1953), Silicate Cement: Method of Mixing in a Closed Container to Prevent Effects of Exposure to Atmosphere, *J. Amer. dent. Ass.*, **46**, 184.
- Handbook of Chemistry and Physics, 1953, p. 1250. Cleveland: Chemical Rubber Publishing Company.
- MATTHEWS, ERNEST (1935), Silicate Cements, *Proc. R. Soc. Med.*, **28**, 742.
- PAFFENBARGER, G. C. (1951), *Silicate Cement and Amalgam Fillings, Current Advances in Dentistry 1950-1951*. Illinois: The University of Illinois Press.
- SCHOONOVER, IRL C., and SOUDER, WILMER (1938), Dental Silicate Cements: Physical and Chemical Properties and a Specification, *J. Amer. dent. Ass.*, **25**, 32.
- PHILLIPS, RALPH W., and SWARTZ, MARJORIE L. (1957), Effect of Certain Restorative Materials on the Solubility of Enamel, *Ibid.*, **54**, 623.
- VOLKER, J., BEKARIS, B., and MELILLO, S. (1944), Some Observations on the Relationship between Plastic Filling Materials and Dental Caries, *Tufts Dental Outlook*, **18**, 4.

BOOK REVIEWS

THE IMPACTED LOWER THIRD MOLAR.

By WILLIAM E. DURBECK, A.B., D.D.S., Consultant, Robert B. Green Memorial Hospital, San Antonio, Texas; Civilian Consultant, Brooke Army Hospital, Ft. Sam Houston, Texas. Second edition. 9×6 in. Pp. 188+x, with 117 illustrations. 1958. London: Henry Kimpton. (Brooklyn, N.Y.: Dental Items of Interest Publishing Co. Inc.) 37s. 6d.

In many books on dental and oral surgery the impacted third molar has its rightful place, but it is unusual for a book to appear which deals with such a tooth *in toto* and that is why it is such a welcome addition to dental literature. This book appeared in its first edition in 1943 and in the second one there is an additional chapter on sub-acute bacterial endocarditis, which emphasizes the necessity of eliciting a complete

history of the patient who is going to have his impacted third molar extracted. Beginning with the chapter on general considerations the author deals with, in the succeeding eight chapters, anatomical considerations, clinical examination and pre-operative treatment, anaesthesia, general roentgen considerations, radiographic examinations, planning the procedure, technique of instrumentation, and post-operative care. In the actual operative technique the reader may find himself at variance with the author's suggestions. Nevertheless, the technique and instrumentation is dealt with on sound basic principles of planning the procedure. The book is fully illustrated, but it is unfortunate that the majority of the illustrations are of poor quality and poorly orientated. This is a minor criticism which should not deter the reader from the value of the book, which should be

in the hands of not only oral surgeons but general dental practitioners as well. This book will well repay careful study and can be heartily recommended.

H. M.

ORAL MICROBIOLOGY AND INFECTIOUS

DISEASE. A Textbook for Students and Practitioners of Dentistry. By GEORGE W. BURNETT, B.A., M.A., D.D.S., Ph.D., F.A.C.D. Lieutenant Colonel, Dental Corps, U.S. Army; and HENRY W. SCHERP, B.S., M.S., Ph.D., Associate Professor of Bacteriology and Immunochemistry, University of Rochester, School of Medicine and Dentistry, Rochester. 10×6½ in. Pp. 589+xvi, with 257 illustrations. 1957. Baltimore: The Williams and Wilkins Co. (London: Baillière, Tindall & Cox Ltd.) 88s.

THIS book may be divided into three parts: the first part, besides including a section on the history of microbiology, covers the cultivation, cytology, and metabolism of bacteria, with an account of sterilization, chemotherapy, and serology. The second part deals with the bacteriology of caries, periodontal disease, pulpal and periapical infections, with chapters

on those systemic diseases caused by micro-organisms and with oral manifestations. Lastly, there are descriptions of individual organisms and their associated diseases.

For those readers who already have a good knowledge of the subject, this very well-illustrated book should be most useful. Quite apart from the lucidity of description, it would be difficult to name another text where the bacteriological aspects of those diseases of immediate interest to the dental surgeon are dealt with in such a competent fashion. However, its suitability for use by undergraduates, except as a reference book, is another matter. The length of the book (589 pages) indicates its detail. An undergraduate, and indeed many postgraduates, would tend to be overwhelmed by the relatively detailed presentation, especially regarding chapters such as that on serology.

It would be fair to say that this is a very good book for a postgraduate specializing in this subject, but is not very suitable as a standard text for the undergraduate dental student in this country.

E. N.

ABSTRACTS FROM OTHER JOURNALS

Ethyl Chloride, Nitrous Oxide, and Oxygen Mixture

In answer to the question "I give a number of dental anaesthetics, chiefly to children, using a Mekesson Apparatus, with vinyl-ether in addition to nitrous oxide and oxygen for prolonged anaesthesia. I would prefer, however, to give ethyl chloride in the nitrous oxide and oxygen mixture, as in many ways this is safer and better than vinyl-ether or trichloroethylene. Is there any apparatus which can be incorporated with a Mekesson for delivering ethyl chloride by drip or by controlled addition to the mixture? Is there any objection to the idea of a nitrous oxide-oxygen mixture with ethyl chloride in an open circuit?"

The following answer is given:

"Ethyl chloride as a supplement to nitrous oxide and oxygen has been in use for a very

long time. Within the last fifteen years serious interest in this combination has been revived. There is no commercially available vaporizer for ethyl chloride for the Mekesson apparatus, nor is there one for such British equipment as the Boyle or Walton. However, all that is necessary is a small hole in the circuit somewhere between the apparatus and patient into which ethyl chloride can be squirted. A drip vaporizer would need special precautions in its design owing to the low boiling-point of ethyl chloride. There is no objection to ethyl chloride being used in an 'open' circuit. Ethyl chloride is widely used, its main danger being overdose. Many anaesthetists, however, particularly in the United States, would not agree with the questioner that ethyl chloride 'is safer and better than vinyl ether or trichloroethylene'."—*Brit. med. J.* (1958), 1, 236.

Fibrous Dysplasia of Maxilla and Mandible

A clinical, radiological, and histological study has been made of 69 cases of fibrous dysplasia of the maxilla and the mandible, seen at the Mayo Clinic.

It is an indolent tumefactive process of fibro-osseous character seen in patients between the ages of 10 and 30 years. It occurs more often in the maxilla of the female. The pathogenesis is obscure, but it is considered to be due to a developmental defect. Many fibro-osseous lesions fall within the scope of fibrous dysplasia, but conditions which can be separated by clinical, laboratory, and roentgenological criteria such as cementoma, osteoid osteoma, epulis with calcification, cherubism, brown tumours of hyperparathyroidism, and Paget's disease should not be classified under fibrous dysplasia. Chondromatous elements are rare in maxillary and mandibular fibrous dysplasia. If present, it is usually malignant. The treatment of choice in fibrous dysplasia of the jaws is conservative surgical intervention. Irradiation is of no value. Approximately 20 per cent of these tumours continue to grow after treatment, except for those radically excised.—ZIMMERMAN, D. C., DAHLIN, D. C., and STAFNE, E. C. (1958), *Oral Surg.*, 2, 55.

Basic Principles of Endodontia and their Application to Selection and Treatment

The following list of ten basic principles applies to every endodontic case: (1) Treatment of pulp-involved teeth is attempted only on those patients interested in retaining their teeth. (2) The patient's health should be sufficiently good to withstand the demands of proper treatment. (3) The tooth must be accessible and anatomically operable. (4) The tooth must be restorable. (5) The tooth must be strategic in relation to the dentition as a whole. (6) Periodontal tissue surrounding the tooth should be within normal limits to ensure adequate healthy support following completion of endodontic treatment. (7) If traumatic occlusion is present it must be capable of being corrected. (8) Canals should be free from debris and dry before being filled. If

surgery does not follow filling of the canals, the canals should be sterile as established by two successive negative cultures. (9) Canals should be sealed hermetically and neither be over-filled nor underfilled. The ideal filling will obliterate the canal at the cemento-dentine junction. (10) All areas denoting pathologic lesions must show evidence of repair on follow-up X-ray examinations.—CRAIN, E. L. (1958), *Northw. Dent.*, 37, No. 1.

Severe Reactions to Antibiotics

The problem of reactions to antibiotics, especially penicillin, is an increasingly grave one. The survey covered a three-year period and included a study of approximately one-third of the general hospital beds available in the United States. A total of 2995 reactions were reported, 80 per cent of which were associated with the use of penicillin. In those cases where the reaction was sufficiently severe to be life threatening in degree, penicillin was a responsible agent in 84 per cent of the cases. Of the 901 penicillin cases 83 terminated fatally, giving a mortality rate of 9 per cent.

It is of interest to note that the number of reactions to penicillin when administered in capsule or oral form was numerically less than when administered in the injectable form.

The lesson to be learnt from this important study is that this wonderful group of life-saving drugs may also be life threatening if the patient has been sensitized to the drug previously. Therefore, they should not be used promiscuously. Instead, they should be used only when there is a true indication that they are of great importance to the patient's successful recovery. When used, the oral form is obviously the safest route of medication and should be used when at all possible. If injection of penicillin is required, adequate emergency 'know-how' drugs and equipment must be immediately available since the typical patient with an anaphylactoid reaction usually dies within a few minutes unless emergency treatment is administered immediately.—HANNY, F. A. (1958) *J. Mich. St. dent. Assoc.*, March, 45.

Class 2 Cavity Preparations for Amalgam

In a survey of 1000 failures of amalgam fillings it is estimated that about 85 per cent were due to improper cavity preparation with poor extension for prevention, the most common error; 60 per cent were due to faulty cavity preparation.

Class 2 cavity undoubtedly offers the greatest opportunity for error in preparation. The occlusal portion must be extended to include all developmental grooves. The pulpal floor should be flat and seated at least a millimetre beyond the dento-enamel junction, the buccal and lingual cavity wall should converge slightly towards the occlusal to give what is termed the inverted truncated cone form, while the caries-free approximal cavity wall should either be perpendicular to the pulpal floor or diverge slightly towards the occlusal in order to preclude the chance of undermining the intact proximal marginal ridge.

Amalgam has a very good crushing strength if there is sufficient bulk to the material, but frail feather-edged margins cannot be tolerated. A bevel must always be avoided on the margins of a cavity preparation for amalgam.

There are five very definite reasons for cutting the cavity walls in the manner described in the text, and these five reasons are: (1) Improved resistance form; (2) Improved retention form; (3) Reduced chance of establishing a feather-edged margin at the occluso-bucco-proximal and occluso-proximal point angles; (4) Conservation of more of the important marginal ridge of the tooth; (5) Placement of the gingivo-bucco-proximal and gingivo-linguo-proximal point angles in areas of relative immunity to caries. When both proximal surfaces of the tooth are involved with caries two class 2 cavities are prepared and are joined on the occlusal surface.

Regardless of the instruments employed, such as ultra-high-speed handpieces and drills, the principles of cavity preparation remain the same. The newly invented high-speed drills should be used with great consciousness and wisdom.—MOSTELLER, J. H. (1958), *Northw. Dent.*, 37, No. 1.

More Aids to Endodontic Practice

The author gives a number of additional aids with the object of saving time and labour, and so making the practice of endodontics more pleasant and comfortable for the operator. Some aids are original. It is recommended that these aids be put into practice as soon as the opportunity permits.—GROSSMAN, L. I. (1958), *Oral Surg.*, 2, 91.

A Precision and Biological Root Canal Filling Technique

A root canal can be underfilled, over-filled, filled exactly flush with the apical foramen, or filled level with the cemento-dental junction. The advantages and disadvantages of each are discussed and the latter type of filling is favoured by the author as it offers the best chance for the apex to seal itself with cementum.

The various methods of filling a root canal are enumerated—namely, Diffusion technique, Impregnation technique, Overfilling technique, Cement technique, and Condensation technique, and these are assessed by comparison with a list of stipulations for an ideal root filling. Objections are raised against each technique and in order to overcome these objections a new technique is proposed.

This consists of reaming only the dentinal part of the root canal. This is carefully done to the greatest diameter feasible, and using a depth gauge. The taper on a selected standard gutta-percha point is reduced by rolling between glass slabs. The point is tried in the canal and is considered satisfactory when it fits firmly in the canal stopping short of the cemento-dental junction by 0.5 mm. The point is then removed, moistened in chloroform to a depth of 0.5 mm., dipped into dentine dust, and pressed firmly into the canal, this time to the proper depth, i.e., to the cemento-dental junction.

The rest of the pulp chamber and canal is filled with a silver cement introduced with smooth branches and packed with thin accessory gutta-percha points. Of 451 root canals filled in this way the periapical tissues failed to regenerate in 5 only.—KUTTLER, YURY (1958), *J. Amer. dent. Assoc.*, 56, 38.

INDEX

VOLUME VIII, Nos. 1—12 (SEPTEMBER, 1957—AUGUST, 1958)

	PAGE		PAGE
A			
ABNORMALITY, growth and the prediction of	220	BENNETT, D. T., and NICHOLS, P. J. R.: Observations on two anaesthetic agents, xylodase paste and xylotox extra paste	204
Abstracts	18, 29, 30, 41, 61, 80, 94, 121, 128, 135, 139, 150, 160, 174, 176, 184, 194, 203, 214, 219, 231, 240, 255, 257, 274, 285, 297, 304, 308, 319, 326, 348, 365, 386, 391	BLACKMAN, SYDNEY: Radiation hazards in odontostomatology	209
Acrylic resin implants, self-curing, tissue reaction to	234	— — Radiographic determination of periapical disease	110
Aetiology of malocclusion	42	Bone implant, histology of	26
Alloy(s), aluminium casting, use of, for constructing denture bases	382	Book Reviews (see Reviews)	
— yellow and white gold and cobalt-chromium, a comparison of	8	BOUCHER, CARL O.: A fundamental approach to the problems of impressions for complete dentures	162
Aluminium casting alloy for constructing denture bases, use of	382	Bridge and crown ceramics and rehabilitation, provisional coverage and splinting procedures in	198
Alveolar process, study of growth in height of: O'Meyer's sign	140	BROADWAY, R. T.: Depression of lower incisors	192
Amalgam restorations, finishing	64	BROWNE, WILLIAM G.: Surgical assistance for the eruption of four anterior maxillary teeth	28
American Dental Society of Europe: Developmental supervision of occlusion, 81; A fundamental approach to the problems of impressions for complete dentures, 162; Practical periodontitis treatment easily applied by the general practitioner, 280; Principles of complete denture prosthesis, 350; Principles of ultrasonic cavity preparation, 122; Provisional coverage and splinting procedures in crown and bridge ceramics and rehabilitation, 198; Silicate cements: How to select and use them, 387; The use of plate appliances in the treatment of periodontal trauma, 12; The use of suggestion and psychosomatics in dentistry, 312		British Society of Periodontology, Proceedings of: Programme, 1957-8, 60; Annual clinical meeting, 1957, 19, 57; Calculus and wear in the teeth of sixth century Jutes, 23; A case of periodontosis in a boy aged $6\frac{1}{2}$ years, 58; A chemical change in cervical dentine in periodontal disease, 19; Chrome-cobalt subperiosteal implant, 59; Current trends in periodontal research, 151; Dental health education in Italy, Switzerland, and Germany, 286; Fundamentals of collagen, 261; Heterogenous bone implant, 58; The histology of a bone implant, 26; Homogenous bone implant, 57; A new method for measuring tooth mobility, 329; Osteoplasty, 59; Periodontal splint, 22; The periodontal tissues of some laboratory animals, 53; Practical periodontics, 215; Recurrent aphthous ulcers, 177; Seasonal incidence of acute gingivitis, 23; Some of the less common periodontal lesions, 23; The staple-type of fixed splint, 22	
Anæsthetic agents, xylodase paste and xylotox extra paste, observations on	204	British Society for the Study of Orthodontics, Transactions of: The aetiology of malocclusion—an assessment, 42; Depression of lower incisors, 192; The eruption of third molars following extraction of second molars, 292; Growth and the prediction of abnormality, 220; An investigation into the effects of pressure on various bony sites in the rabbit, 295; A mobile orthodontic chairside desk, 256; Report on the edgewise appliance, 86; A study of the growth in height of the alveolar process: O'Meyer's sign, 140; The treatment of Angle's Class II, division 1 and Class II, division 2 in identical twins, 185; Treatment of the unerupted maxillary canine, 245; The use of contoured canine orthodontic bands, 191	
Analysis of one hundred cases of dry socket	275		
ANDERSON, J. N.: The dimensional stability of three silicone-base impression materials	368		
— — Dimensions of cast palatal and lingual bars	270		
ANGELMAN, JOSEPH: A new apron spring	51		
Angle's Class II, division 1 and Class II, division 2 in identical twins, treatment of	185		
Animals, laboratory, the periodontal tissues of some	53		
Anterior maxillary teeth, surgical assistance for eruption of four	28		
Aphthous ulcers, recurrent	177		
Appliance, edgewise, report on	86		
Apron spring, a new	51		
ASTBURY, W. T.: Fundamentals of collagen	261		
B			
BALLARD, C. F.: The aetiology of malocclusion—an assessment	42		

British Society for the Study of Prosthetic Dentistry:

The design of lower free-end saddle dentures, 322; Dimensions of cast palatal and lingual bars, 270; The non-metallic denture base—recent developments, 73; Stresses in denture bases, 37

C

CAMPBELL, J. MENZIES: The Crawcours	108
Canine orthodontic bands, contoured, use of	191
— unerupted maxillary, treatment of	245
Cavity preparation, pain and the innervation of dentine in	171
— ultrasonic, principles of	122
Cements, silicate: How to select and use them	387
Cephalometric analysis suitable for the different constitutional types of head, new method of	358
Ceramics and rehabilitation, crown and bridge, provisional coverage and splinting procedures in	198
CLARKE, J.: see HAMPSON, E. L., jt. auth.	
Cobalt-chromium alloys and yellow and white gold alloys, a comparison of	8
Collagen, fundamentals of	261
COLLINGE, J. S. HILTON: An interesting dentigerous cyst	340
COOK, H. P.: Facial fibrous dysplasia	105
Copper dies, electro-formed, from impressions in elastic materials	70
Courses for dental technicians	30
Crawcours, the	108
CRIPPS, S.: Practical periodontics	215
CROSS, W. G.: The histology of a bone implant	26
Crown and bridge ceramics and rehabilitation, provisional coverage and splinting procedures in	198
— forms, aberrant, a compression odontome and its possible relationship to	309
— post-retained	130
Cyst, dentigerous, an interesting	340

D

DENTAL health education in Italy, Switzerland, and Germany	286
— laboratory, a protective shield for use in	175
— practice, hypnosis and its limitations in	258
— tissues, hard, cutting of, advances in	98
Dentigerous cyst, an interesting	340
Dentine, pain and the innervation of, in cavity preparation	171
Dentistry, the use of suggestion and psychosomatics in	312
Dentition, effect of thumb-sucking on	300
Denture(s) base(s), aluminium casting alloy for construction of, use of	382
— non-metallic—recent developments	73
— stresses in	37
— complete, a fundamental approach to the problems of impressions for	162
— lower free-end saddle, design of	322
— prosthesis, complete, principles of	350
Desk, orthodontic chairside, mobile	256
Dies, copper, electro-formed, from impressions in elastic materials	70

PAGE

PAGE

Disease, periapical, radiographic determination of	110
DURKACZ, K. P., and MGLEJ, S. F.: The value of ultra-violet light in macroscopic examination of teeth	136
Dysplasia, fibrous, facial	105

E

EDGEWISE appliance, report on	86
Editorial 1, 31, 63, 97, 129, 161, 197, 233, 269, 299, 335, 367	
Elastic materials, electro-formed copper dies from impressions in	70
Electromyography relative to stomatognathic system	32
Enamel surface of human tooth, integument of	341, 373
ENGLISH, JAMES A.: Current trends in periodontal research	151
Eruption of four anterior maxillary teeth, surgical assistance for	28
— third molars following extraction of second molars	292
Extraction of second molars, eruption of third molars following	292

F

FACIAL fibrous dysplasia	105
— growth, studies in	326
FARMER, E. DESMOND: Recurrent aphthous ulcers	177
FEARNHEAD, R. W.: Pain and the innervation of dentine in cavity preparation	171
FERGUSON, H. W.: see HITCHIN, A. D., jt. auth.	
Fibrous dysplasia, facial	105
FISHER, D. KEIR: Hypnosis and its limitations in dental practice	258
FRANKS, ARNOLD S. T.: Electromyography relative to the stomatognathic system: a review	32
Furnace, gas-heated vacuum, for porcelain	305

G

GERMANY, Italy, and Switzerland, dental health education in	286
Gingival trough, implant materials and	2
Glasgow Odontological Society: Hypnosis and its limitations in dental practice, 258; Surgical assistance for the eruption of four anterior maxillary teeth, 28	
Gold alloys, yellow and white, and cobalt-chromium alloys, a comparison of	8
GOOSE, D. H.: Dental health education in Italy, Switzerland, and Germany	286
GRATZINGER, MAX: The use of plate appliances in the treatment of periodontal trauma	12
Growth, facial, studies in	327

H

HALDEN, J. R.: Report on the edgewise appliance	86
HAMPSON, E. L., and CLARKE, J.: The post-retained crown	130

	PAGE		PAGE
Head, different constitutional types of, new method of cephalometric analysis suitable for	358	LUCCHESI, PAOLO: see MAJ, GIORGIO, jt. auth.	
HENRY, R. G.: The effect of thumb-sucking on the dentition	300	LUZI, CESARE: see MAJ, GIORGIO, jt. auth.	
HEYLINGS, R. TAYLOR: The periodontal tissues of some laboratory animals	53	LYON, F. F.: Gas-heated vacuum furnace for porcelain	305
Histology of bone implant	26		
HITCHIN, A. D., and FERGUSON, H. W.: A compression odontome and its possible relationship to aberrant crown forms	309	M	
HUDSON, W. CAMPBELL: Provisional coverage and splinting procedures in crown and bridge ceramics and rehabilitation	198	McCAGIE, J. N. W.: A case of re-implantation of teeth after five days	320
Hypnosis and its limitations in dental practice	258	Macroscopic examination of teeth, the value of ultra-violet light in	136
		MAJ, GIORGIO, LUZI, CESARE, and LUCCHESI, PAOLO: A new method of cephalometric analysis suitable for the different constitutional types of head	358
I		Malocclusion, aetiology of	42
IMPLANT(s), bone, histology of	26	Maxillary canine, unerupted treatment of	245
— materials and gingival trough	2	— teeth, four anterior, surgical assistance for eruption of	28
— self-curing acrylic resin, tissue reaction to	234	MESSING, J. J.: Linings and their manipulation	336
Impression(s) for complete dentures, a fundamental approach to the problems of	162	MCLEJ, S. F.: see DURKACZ, L. P., jt. auth.	
— materials, dimensional stability of three silicone-base	368	Molars, deciduous, role of retained upper second in the causation of post-normal occlusion	207
Incisors, lower, depression of	192	— third, eruption of, following extraction of second	292
Italy, Switzerland, and Germany, dental health education in	286	MORRANT, GUY A.: Principles of ultrasonic cavity preparation	122
J		N	
JAMIESON, CHARLES H.: Principles of complete denture prosthesis	350	NEILL, D. J.: The design of lower free-end saddle dentures	322
JOEL, A. A.: A new method for measuring tooth mobility	329	— — Use of aluminium casting alloy for constructing denture bases	382
JOHNSON, W.: A comparison of cobalt-chromium alloys and yellow and white gold alloys	8	New Materials and Appliances: Compressed-air handpiece, 159; Copper-forming dental unit, 159; Dental furnace, 104	
		NICHOLLS, E.: Sectional root filling with silver points	241
K		NICHOLS, P. J. R.: see BENNETT, D. T., jt. auth.	
KANTOROWICZ, G. F.: Finishing amalgam restorations	64	Norwegian Institute of Dental Research: Tissue reaction to self-curing acrylic resin implants	234
KETTLE, M. A.: Treatment of the unerupted maxillary canine	245		
KNOWLES, C. C.: The role of retained upper second deciduous molars in the causation of post-normal occlusion	207	O	
		Obituary Notice: J. Forbes Webster	334
L		Occlusion, developmental supervision of	81
LABORATORY animals, periodontal tissues of some	53	— post-normal, role of retained upper second deciduous molars in causation of	207
— dental, a protective shield for use in	175	Odontome, compression, and its possible relationship to aberrant crown forms	309
LAMMIE, G. A.: Advances in the cutting of hard dental tissues	98	Odonto-stomatology, radiation hazards in	209
LAMONS, FRANK F.: Developmental supervision of occlusion	81	OLDFIELD, H. M.: see STORER, ROY, jt. auth.	
LEECH, H. L.: The treatment of Angle's Class II, division 1 and Class II, division 2 in identical twins	185	O'MEYER, R. X.: A study of the growth in height of the alveolar process: O'Meyer's sign	140
LEHNER, T.: Analysis of one hundred cases of dry socket	275	Orthodontic(s) bands, contoured canine, use of	191
Letters to the Editor	60, 125, 160, 195, 279	— British Society for the Study of: (see British Society for the Study of Orthodontics)	
Lingual and palatal bars, cast, dimensions of	270	— desk, mobile chairside	256
Linings and their manipulation	336		
LOE, HARALD: see WAERHAUG, JENS, jt. auth.		P	
Lower free-end saddle dentures, design of	322	PAFFENBARGER, GEORGE C.: Silicate cements: How to select and use them	387
		Palatal and lingual bars, cast, dimensions of	270
		PEARSON, S. L.: Electro-formed copper dies from impressions in elastic materials	70

	PAGE		PAGE
Periapical disease, radiographic determination of -	110	Reviews:—continued	
Periodontal research, current trends in -	151	<i>Oral histology and embryology</i> (ed. Balint J. Orban) -	96
— trauma, use of plate appliances in treatment of	12	<i>Oral hygiene</i> (Russell W. Bunting) -	158
Periodontics, practical -	215	<i>Oral medicine</i> (Lester W. Burket) -	267
Periodontitis, practical treatment easily applied		<i>Pediatric dentistry</i> (M. Michael Cohen) -	7
by the general practitioner -	280	<i>Phantomkurs der Konservierenden Zahnheilkunde</i>	
Periodontology, British Society of: (see British		(C. H. Rathner) -	7
Society of Periodontology)		<i>Pharmacologie et pharmacotherapie dentaire</i>	
PERSTON, C. M.: An investigation into the effects of		(O. Noeppel) -	349
pressure on various bony sites in the rabbit	295	<i>Regulations of dental practice</i> -	158
Plate appliances in treatment of periodontal		<i>Teething troubles</i> (Matthew Finch) -	230
trauma, use of -	12	<i>The third set</i> (Matthew Finch) -	230
Porcelain, gas-heated vacuum furnace for -	305	<i>Xylocaine in anesthesia</i> -	231
Post-normal occlusion, role of retained upper		RIDLEY, DORIS R.: The use of contoured canine	
second deciduous molars in causation of -	207	orthodontic bands -	191
Post-retained crown -	130	Root filling, sectional, with silver points -	241
Practical periodontics -	215	ROWE, F. D.: A mobile orthodontic chairside desk	256
— periodontitis, treatment easily applied by		Ruspini, Chevalier Bartholomew -	298
general practitioner -	280		
Psychosomatics, use of suggestion and, in dentistry	312		
		S	
R		SADDLE dentures, design of lower free-end -	322
RABBIT, an investigation into the effects of pressure		SCOTT, JAMES H.: Studies in facial growth -	326
on various bony sites in -	295	SHAW, S. IRWIN: The use of suggestion and psycho-	
Radiation hazards in odonto-stomatology -	209	somatics in dentistry -	312
Radiographic determination of periapical disease -	110	Shield, protective, for use in dental laboratory -	175
Rehabilitation and ceramics, crown and bridge,		Silicate cements: How to select and use them -	387
provisional coverage and splinting		Silicone-base impression materials, dimensional	
procedures in -	198	stability of three -	368
Re-implantation of teeth after five days -	320	Silver points, sectional root filling with -	241
Research, periodontal, current trends in -	151	SMITH, D. C.: The non-metallic denture base—	
Resin implants, self-curing acrylic, tissue reaction		recent developments -	73
to -	234	SMITH, D. I.: The eruption of third molars following	
Restorations, amalgam, finishing -	64	extraction of second molars -	292
Reviews:—		Society for the Advancement of Anesthesia in	
<i>The Aleut dentition</i> (Coenraad F. A. Moorrees) -	332	Dentistry -	85
<i>Calcium metabolism</i> (J. T. Irving) -	127	Society of Medical Officers of Health: Radiation	
<i>The century of the surgeon</i> (Jürgen Thorwald) -	196	hazards in odonto-stomatology -	209
<i>Cleft lip and palate</i> (W. G. Holdsworth) -	126	Socket, dry, analysis of one hundred cases of -	275
<i>Clinical applications of hypnosis in dentistry</i>		Spring, apron, a new -	51
(S. Irwin Shaw) -	333	Stomatognathic system, electromyography relative	
<i>Clinical orthodontics</i> (Bera Fischer) -	229	to -	32
<i>Clinical pedodontics</i> (Sidney B. Finn) -	230	STORER, ROY, and OLDFIELD, H. M.: A protective	
<i>Clinical procedures in occlusal rehabilitation</i>		shield for use in the dental laboratory -	175
(S. Charles Brecker) -	349	Stresses in denture bases -	37
<i>Dental clinics of North America</i> (ed. Hamilton		Supervision, developmental, of occlusion -	81
B. G. Robinson and Robert G. Kesel) -	296	Surgical assistance for eruption of four anterior	
<i>Dental clinics of North America</i> (ed. F. Darl		maxillary teeth -	28
Ostrander) -	333	Switzerland, Italy, and Germany, dental health	
<i>Dental health education</i> (Frances A. Stoll) -	126	education in -	286
<i>Dental practitioners' formulary, 1957</i> -	296		
<i>The design and construction of removable ortho-</i>		T	
<i>dontic appliances</i> (C. P. Adams) -	268	TANNER, J. M.: Growth and the prediction of	
<i>The fight for fluoridation</i> (Donald R. McNeil) -	230	abnormality -	220
<i>The impacted lower third molar</i> (William E.		Teeth, four anterior maxillary, surgical assistance for	
Durbeck) -	390	eruption of -	28
<i>Implant dentures</i> (Aaron Gershkoff) -	267	— human, integument of the enamel surface of	
<i>Local anesthesia and pain control in dental practice</i>		341, 373	
(Leonard M. Monheim) -	127	— macroscopic examination of, value of ultra-	
<i>A manual of orthodontics</i> (Hallam Gresham) -	332	violet light in -	136
<i>A manual of stomatology</i> (J. P. Walsh) -	95	— re-implantation of, after five days -	320
<i>Oral diagnosis and treatment</i> (Samuel Charles		Thumb-sucking, effect of on dentition -	300
Miller) -	95	Tissue(s), hard dental, cutting of, advances in -	98
<i>Oral and facial cancer</i> (Bernard G. Sarnat and		— periodontal, of some laboratory animals -	53
Isaac Schour) -	195	— reaction to self-curing acrylic resin implants -	234

	PAGE		PAGE
Tooth mobility, new method for measuring -	329	V	
TRAININ, BORIS: Implant materials and the gingival trough -	2	VACUUM furnace, gas-heated, for porcelain -	305
Trauma, periodontal, use of plate appliances in treatment of -	12		
Treatment of Angle's Class II, division 1 and Class II, division 2 in identical twins -	185	W	
— periodontal trauma, use of plate appliances in -	12	WAERHAUG, JENS, and LOE, HARALD: Tissue reaction to self-curing acrylic resin implants -	234
— unerupted maxillary canine -	245	WAIN, E. A.: Stresses in denture bases -	37
TURNER, E. P.: The integument of the enamel surface of the human tooth -	341, 373	WIESSTIEN, L. M.: Practical periodontitis treatment easily applied by the general practitioner -	280
Twins, identical, treatment of Angle's Class II, division 1 and Class II, division 2 in -	185		
		X	
U			
ULCERS, aphthous, recurrent -	177	XYLODASE paste and xylotox extra paste, observations on -	204
Ultrasonic cavity preparation, principles of -	122	Xylotox extra paste and xylodase paste, observations on -	204
Ultra-violet light in macroscopic examination of teeth, value of -	136		

THE DENTAL PRACTITIONER and DENTAL RECORD

SEP 16 1958

VOL. VIII, NO. 12

DENTISTRY
LIBRARY

AUGUST, 1958

Including the official reports of the British Society of Periodontology, the British Society for the Study of Orthodontics, the European Orthodontic Society, the Glasgow Odontological Society, the Liverpool and District Odontological Society, the North Staffordshire Society of Dental Surgeons, the Odontochirurgical Society of Scotland, and the Dental and Medical Society for the Study of Hypnosis

CONTENTS

~~DOES NOT CIRCULATE~~

Editorial : Fluoridation and Andover

The Dimensional Stability of Three Silicone-base Impression Materials
J. N. ANDERSON, M.D.S., L.D.S. (Sheff.)

The Integument of the Enamel Surface of the Human Tooth (*continued*)
E. P. TURNER, M.Sc., L.D.S. R.C.S. (Edin.)

Use of an Aluminium Casting Alloy for Constructing Denture Bases
D. J. NEILL, D.F.C., F.D.S.


Silicate Cements : How to Select and Use Them
GEORGE C. PAFFENBARGER, D.D.S., D.Sc.

Book Reviews

Abstracts from Other Journals

JOHN WRIGHT & SONS LTD., BATH ROAD, BRISTOL 4

THREE SHILLINGS AND SIXPENCE




...as possible.
 2725-2845 Local Government Superannuation Scheme.—Reply by 31st May, enclosing two references, to: The Principal, Arlington Manor, Newbury, Berks.

STOLEN. Entire contents of safe: Jewels, Cash, Bonds, etc. including Upper Denture fitted with Classic 'S' Teeth. If burglar will return denture which was specially made for owner, guarantee no police action in respect of other items.

MEDICAL SECRETARY required for work in the Radiotherapy Department. Grade of the N.H.S. scales.—Apply in writing, to: previous experience.

A
 asst
 recd
 Aut
 31
 Co
 Er

AT LAST...



a safe, simple and positive treatment for

Aphthous Ulcers

Now, at last, in Corlan Pellets you have a safe, simple and positive treatment for aphthous ulceration*—a condition that hitherto has failed to respond to a wide variety of medicaments.

Corlan is a special water-soluble form of hydrocortisone presented in pellet form in a base designed to dissolve slowly in the saliva.

CORLAN PELLETS

TRADE MARK


*"The method of treatment we have described is not only effective in the great majority of patients but is also simple and safe. A tablet is easily held in the mouth and will rest in the gingival sulcus, where aphthous ulcers most commonly occur. Its slow dissolution ensures that the ulcer is bathed in saliva charged with hydrocortisone for a rather long period."

Brit. Med. J., 1958, **1**, 603

*Each pellet contains 2.5 mg.
hydrocortisone in the form of the
water-soluble ester, hydrocortisone
hemisuccinate sodium.
Pack: Tube of 20.*



GLAXO LABORATORIES LTD., GREENFORD, MIDDLESEX. BYR0n 3434



Kemdent

DISCS and MANDRELS

made for each other . . .

Secure accurate location of disc on mandrel every time, firm anchorage and steady support by using the specially matched KEMDENT Type 'B' Discs with the new KEMDENT Type 'B' Mandrel (Regd. Design). This is a precision job with an easy, snap-on action. Type 'B' Mandrels are available for No. 7 and No. 2 R.A. Handpieces.

A DISC TRAY CASE

- *A neat, handy*
- *case in black*
- *leatherette with*
- *nine compart-*
- *ments to hold an*
- *assortment of*
- *discs and a*
- *reference chart.*
- *Supplied empty*
- *or filled with 600*
- *assorted Type 'B'*
- *Discs.*



Kemdent Standard Pin Hole Discs (Type A) are available in the usual Grits

Available from usual Dental Depots

Manufactured by

ASSOCIATED DENTAL PRODUCTS LIMITED

Purton, Swindon, Wilts. & London, W.1



Safety First !!!!!

Nearly a quarter of a century of proven service places ANADIN in the forefront of modern analgesics. Its unparalleled record of unobtrusive efficiency even in patients with established systemic disease has given it a unique place in dental practice.

ANADIN is a safe and efficient combination of drugs capable of providing speedy relief of post-operative pain. The complete absence of side effects makes it the analgesic of choice following oral surgery.

Two tablets at the first sign of pain ensure swift and effective relief with just the right amount of mental stimulation to banish worry and fear.

Anadin

Trade Mark
TABLETS

INTERNATIONAL CHEMICAL COMPANY LTD.,
CHENIES STREET, LONDON, W.C.1

ARE YOU A VICTIM OF...



CAPITAL PUNISHMENT

Today there is
no longer a necessity
for Dentists in need
of new equipment to
break into
capital resources—
we have made it
possible to equip a
surgery
without initial cost,
by the introduction of
our "Unique Rental Plan.

Please send or telephone for further details

LANGHAM 2011

DENTAL RENTALS LTD

in association with

THE DENTAL MANUFACTURING CO. LTD., 97 GREAT PORTLAND STREET, LONDON, W.1

xylodase

REGD.

in dental surgery

'Xylocaine' and hyaluronidase (the spreading factor) are compounded in a special base to give XYLODASE its valuable surface anaesthetic properties. Applied to the gums or to the inside of tooth cavities XYLODASE helps the apprehensive patient. The usual delay while local anaesthetic ointments take effect is eliminated.

In tubes of 15 grammes

XYLODASE' is a registered trade mark, the property of Duncan, Flockhart & Co. Ltd. of Edinburgh.

XYLOCAINE' is a registered trade mark, the property of A. B. Astra of Sweden and used by Duncan, Flockhart & Co. Ltd. under licence.

XYLOCAINE' Dental Cartridges, now distributed in the United Kingdom by Astrapharm Ltd., are available from your usual Dental Depot.



DUNCAN FLOCKHART OF EDINBURGH

WHAT IS THE VALUE OF CLINICAL RESEARCH

CLINICAL research, if conducted on a sufficiently wide basis and under strict, scientific control, is the only measure by which a Medical or Dental Practitioner can evaluate a new drug or dentifrice.

Indeed the only recent introduction, which has been proved by seriously conducted clinical tests on patients of all ages, to bring a significant reduction in caries incidence, is the one High-Urea dentifrice—Amm-i-dent.

Amm-i-dent, in paste and powder form, combines carbamide (Synthetic Urea) with diammonium phosphate. This combination not only raises the pH of the mouth after use but lowers the Lactobacillus count proportionately.

Its remarkable value in reducing caries incidence has been proved in every test to which it has been submitted. These are three typical examples.

DURATION OF STUDY	NUMBER OF PATIENTS			CARIES RATE		% Reduction of Caries Incidence by High-Urea Ammoniated Dentifrice
	Total	Control	Test	Control	Test	
4-year study complete report (2)	185	75	110	2.33	1.31	43.6%
3-year study interim report (5)	120	31	89	2.19	1.08	50.9%
2-year study interim report (1)	60	30	30	1.60	0.96	39.6%

References. 1. Gale, J. A., *Dent. Record*, 71, 15, 1951. 2. Henschel, C. J. and Lieber, L., *Oral Surg. Oral Med. and Oral Path.*, 5, 155, 1952. 3. Lefkowitz, W. and Venti, V. I., *Oral Surg. Oral Med. and Oral Path.*, 4, 1576, 1951.

Withal, Amm-i-dent presents no "patient-resistance". It has a very smooth texture, it cleans the teeth beautifully white and it has a most refreshing flavour.

Professional samples and full literature will gladly be sent by the manufacturers on receipt of a card addressed to Professional Dept., Stafford-Miller Ltd., 166 Great North Road, Hatfield, Herts.

It's just typical of

ATTENBOROUGH'S

SERVICE

*Which, over the years
has been faithfully pursued.*

YOU CAN BUILD
A PRACTICE ON IT



VISCOSA HOUSE

C. & L. E. ATTENBOROUGH LTD.

VISCOSA HOUSE • GEORGE STREET • NOTTINGHAM

Telephone: Nottingham 40374

Telegrams: Lateral, Nottingham

O'Hara Limited

Telephone—41497

Managing Director: S. C. Jeffery

CHROME COBALT CASTINGS BY THE WISIL TECHNIQUE

British Craftsmanship backed by "KRUPP" experience.
The Chrome Cobalt base with clasps you can adjust.
Discounts to bona-fide Laboratories.

PRECISION—

PLUS THE PERSONAL TOUCH

Convent Street,
Parliament Street,
Nottingham.

N.W. ↙N↗ N.E.

PADddington 9635**ASHLEY**PADddington 4793

DENTAL LABORATORIES

(Technical Advisers to the Dental Manufacturing Co. Ltd.)

FOR HIGH-CLASS PROSTHETICS

5 Circus Mews, Enford Street, W.1

W ←----- Daily Collection -----→ E

THE DESIGN AND CONSTRUCTION OF REMOVABLE ORTHODONTIC APPLIANCES

By C. P. ADAMS, B.D.S., F.D.S.

Price 21s., postage 1s.

Second Edition. Fully Cloth Bound. 9¾ × 7¼ in. 120 pp. 173 Illustrations

An attempt to clarify and rationalize the problems of the construction of removable orthodontic appliances; the book is not in any sense a guide to diagnosis or treatment planning. The basic principles underlying the action of all types of appliance, whether fixed or removable, are briefly discussed so as to give a picture of the mechanical possibilities inherent in different techniques.

"This second edition is most highly recommended, not only to orthodontists but also to general practitioners who wish to do minor tooth movement in conjunction with the restoration of missing teeth and for periodontal therapy."—*Oral Surgery, Oral Medicine, and Oral Pathology*.

"A thoroughly useful manual . . . always convincing, and moderately priced; it is not surprising that there has been demand for a second edition."—*Journal of the Royal Naval Medical Service*.

"The illustrations are of the highest standard and this book can be recommended to the dental student, the general practitioner, and the orthodontic specialist."—*British Dental Journal*.

Bristol: JOHN WRIGHT & SONS LTD.

CLASSIFIED ADVERTISEMENTS

All Classifications: 4d. per word (minimum 10s.); Box No. 6d. extra. Compound words count as two words; groups of figures and initials as one word. Series discounts apply as on displayed advertisements.

SITUATIONS VACANT

ST. BARTHOLOMEW'S HOSPITAL, E.C.1. Applications are invited for the post of **Registrar in the Dental Department**, tenable from October, 1, 1958, for a period of two years, subject to re-election at the end of the first year. Duties include every branch of dental and oral surgery, including orthodontics and prosthetics. Preference will be given to candidates holding a dental degree or higher qualifications.

Applications, with the names of two referees, should be submitted to the undersigned within the next fourteen days.

C. C. CARUS-WILSON, *Clerk to the Governors.*

ST. BARTHOLOMEW'S HOSPITAL, E.C.1. Applications are invited for the post of **resident Dental House Surgeon** for a period of six months from a date to be arranged this summer. The appointment is recognized by the Royal College of Surgeons for the purposes of the Fellowship in Dental Surgery, and the holder will gain experience in all branches of dental and oral surgery.

Applications, with the names of two referees, should be submitted to the undersigned as soon as possible.

C. C. CARUS-WILSON, *Clerk to the Governors.*

SITUATION WANTED

DENTIST'S or DOCTOR'S RECEPTIONIST. Lady 21 years of age, 5'8", born and educated London, with dental reception experience in Johannesburg, excellent references, speaks French, seeks similar post Knightsbridge or West End. Miss E. Edgar, 7 Wellington Court, Knightsbridge, S.W.1.

COURSES

H.D.D. (Glasgow), F.D.S. R.C.S. and F.D.S. (Edinburgh), L.D.S., and all other Dental Examinations. Postal Courses for all the above examinations can be commenced at any time. For full details apply: The Secretary, Medical Correspondence College, 19, Welbeck Street, London, W.1.

MISCELLANEOUS

AUTHORS invited to submit MSS all types (including poems) for book publication. Reasonable terms. Stockwell Ltd., Ilfracombe. (Estd. 1898).

MERCURY. We supply clean Mercury (Lab. Grade), price 25s. per lb. All distilled grades stocked. Generous allowance for dirty Mercury. We also buy scrap Dental Amalgam, Gold, Silver and Platinum (in any form) at top market prices. BELGRAVE LTD. (DP), 5 Belgrave Gardens, London, N.W.8. (MAI. 7513). Open Saturdays.

ORTHODONTIC TECHNICIAN offers postal service to practitioners requiring removable appliances of highest standard. **50% discount on first trial appliance.** Write for price list to Box No. 158, DENTAL PRACTITIONER & DENTAL RECORD.

THE DENTIST'S CASH BOOK. Designed by a qualified Accountant for the Profession. The simplest method of recording data necessary to prepare the income and expenditure account which will satisfy Inspectors of Taxes, with full instructions for use. Price 25s. 6d., including tax. The Charnwood Publishing Co. Ltd., Coalville, Leicester.

TO ALL ORTHODONTISTS. Save your valuable time by using **Contoured Orthodontic Bands**, used by the leading Dental Hospitals. Write to Paine Engineering Developments Ltd., Fernhurst, near Haslemere, Surrey.

ENGRAVED BRASS, AND
OXYDISSED BRONZE
PROFESSIONAL

NAME PLATES

With
Permanent
Ceramic
Cream
Enamel
Lettering

Full Size Proof
submitted on receipt of
required words. Leaflet Post Free

G. MAILE & SON LIMITED, Engravers
367 Euston Road, London, N.W.1 Phone: Euston 2938

CHROME COBALT VIRILUM

AND ALL OTHER DENTAL PROSTHETICS

send to

The Laboratory originally appointed
by the Virilium Company as its key
laboratory in the South-West

THE EXETER DENTAL LABORATORY LTD.

24 EAST SOUTHERNHAY
EXETER. Telephone 72606

also at

40 WORRAL ROAD, BRISTOL, 8

**We are as near as
your post box**

FIRM SUCTION WITHOUT IRRITATION



However fine a specimen of prosthetic art you may have constructed, there are times when Corega will prove invaluable. The new denture case, the highly nervous patient, the denture-sore mouth—these and similar instances are indications for the use of Corega. A sprinkle of Powder on the plate provides a suction bond which gives perfect adhesion and enables the inexperienced patient to talk, laugh and eat with complete confidence and comfort the first day. It helps the patient to obtain muscular control of the denture and grow quickly accustomed to its presence.

*Please send for samples which will be sent to you as
always—promptly and without charge*

CORE-GA

PROMOTES DENTURE COMFORT

COREGA CHEMICAL CO., 166 Great North Road, Hatfield, Herts.

Printed and Published by John Wright & Sons Ltd., at The Stonebridge Press, Bath Road, Bristol 4.

8693

